

APPROXIMATION THEOREMS FOR SPACES OF LOCALITIES

SYLVY ANSCOMBE, PHILIP DITTMANN, AND ARNO FEHM

ABSTRACT. The classical Artin–Whaples approximation theorem allows to simultaneously approximate finitely many different elements of a field with respect to finitely many pairwise inequivalent absolute values. Several variants and generalizations exist, for example for finitely many (Krull) valuations, where one usually requires that these are independent, i.e. induce different topologies on the field. Ribenboim proved a generalization for finitely many valuations where the condition of independence is relaxed for a natural compatibility condition, and Ershov proved a statement about simultaneously approximating finitely many different elements with respect to finitely many possibly infinite sets of pairwise independent valuations.

We prove approximation theorems for infinite sets of valuations and orderings without requiring pairwise independence.

1. INTRODUCTION

We fix a field K , elements $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$ and start by recalling the classical approximation theorem for absolute values:

Theorem 1.1 (Artin–Whaples 1945¹). *Let $|\cdot|_1, \dots, |\cdot|_n$ be nontrivial absolute values on K .*

(I) *Assume that $|\cdot|_1, \dots, |\cdot|_n$ are pairwise inequivalent.*

Then there exists $x \in K$ with

$$|x - x_i|_i < |z_i|_i \text{ for } i = 1, \dots, n.$$

Since the non-archimedean absolute values correspond to Krull valuations of rank 1, the following theorem is a generalization in the non-archimedean case:

Theorem 1.2 (Bourbaki²). *Let v_1, \dots, v_n be nontrivial valuations³ on K .*

(I) *Assume that v_1, \dots, v_n are pairwise independent⁴.*

Then there exists $x \in K$ with

$$v_i(x - x_i) > v_i(z_i) \text{ for } i = 1, \dots, n.$$

In the literature, one can find three possible directions of generalizing Theorem 1.2:

Firstly, one can unify Theorems 1.2 and 1.1 to approximate with respect to finitely many pairwise independent valuations and absolute values, like in [PZ87, (4.2) Corollary] or [War89, Cor. 27.14]. This also includes approximation with respect to (not necessarily archimedean) orderings on K , see Theorem 2.8 below.

Secondly, one can relax the condition of pairwise independence in Theorem 1.2, at the expense of introducing a compatibility condition, as done by Nagata [Nag53] and Ribenboim [Rib57].

Theorem 1.3 (Ribenboim 1957⁵). *Let v_1, \dots, v_n be valuations on K .*

(I) *Assume that v_1, \dots, v_n are pairwise incomparable and that if w is a common coarsening of v_i and v_j , $i \neq j$, then $w(x_i - x_j) \geq w(z_i) = w(z_j)$.*

January 10, 2019.

¹Published in [AW45]. See also the historical remarks in [Roq02, §4.2.1]. See also [Lan02, XII.1.2], [EP05, 1.1.3].

²This attribution is taken from [Rib57]. See also [Bou89, VI.7.2 Theorem 1], [EP05, 2.4.1], [Efr06, 10.1.7].

³The term *valuation* in this work always refers to Krull valuations, with value group written additively.

⁴That is, they induce distinct topologies on K , or, equivalently, they have no nontrivial common coarsening.

⁵This is a mild reformulation of [Rib57, Théorème 5], see also [Rib68, p. 136 Théorème 3].

Then there exists $x \in K$ with

$$v_i(x - x_i) > v_i(z_i) \text{ for } i = 1, \dots, n.$$

Thirdly, and most pertinently for us, one can find approximation theorems for possibly infinite subsets of the spaces $S_{\text{ord}}(K)$ of orderings or $S_{\text{val}}(K)$ of valuations on K , under some assumptions on (U)niformity, (T)opology and (I)ndependence (our presentation). The first occurrence of such a ‘block approximation’ theorem seems to be [Ers82, Theorem 1], but there, like in many subsequent works including [Pre85, FHV94, Dar00, Ers06, HJP13], these results apply only to very special fields satisfying some geometric local-global principle, an example of which is the following:

Theorem 1.4 (Prestel 1985⁶). *Let $S_1, \dots, S_n \subseteq S_{\text{ord}}(K)$ pairwise disjoint.*

- (T) *Assume that each S_i is compact in the Harrison topology on $S_{\text{ord}}(K)$ (cf. Section 2).*
- (I) *Assume that K is pseudo-real closed (PRC), that is, every geometrically integral K -variety that has rational points over every real closure of K has a K -rational point.⁷*

Then there exists $x \in K$ with

$$(x - x_i)^2 < z_i^2 \text{ for all orderings in } S_i, \text{ for } i = 1, \dots, n.$$

However, for valuations, Darnière [Dar98] and Ershov [Ers01] proved results for general fields.

Theorem 1.5 (Ershov 2001⁸). *Let $S_1, \dots, S_n \subseteq S_{\text{val}}(K)$ pairwise disjoint.*

- (U) *Assume that there exists a common uniformizer $\pi \in K$ of all $v \in S_1 \cup \dots \cup S_n$.*
- (T) *Assume that each S_i is compact⁹ in the Zariski topology on $S_{\text{val}}(K)$ (cf. Section 2).*
- (I) *Assume that the elements of $S_1 \cup \dots \cup S_n$ are pairwise independent.*

Then there exists $x \in K$ with

$$v(x - x_i) \geq v(z_i) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

The aim of this paper is to prove approximation theorems that generalize Theorem 1.2 in these directions simultaneously; they apply to infinite sets of valuations and orderings, without assuming pairwise independence or the presence of a local-global principle. Our main result in this direction is Theorem 4.6, which is in particular a common generalization of Theorem 1.4 and Theorem 1.5. Instead of attempting to explain the technical assumptions of that main theorem here, we now simply quote three instances of it:

Theorem 1.6. *Let $S_1, \dots, S_n \subseteq S_{\text{val}}(K)$ pairwise disjoint.*

- (U) *Assume that there exists a common uniformizer $\pi \in K$ of all $v \in S_1 \cup \dots \cup S_n$.*
- (T) *Assume that each S_i is compact in the Zariski topology on $S_{\text{val}}(K)$.*
- (I) *Assume that for any valuation w on K with a refinement in some S_i and a refinement in some S_j , we have $w(x_i - x_j) \geq w(z_i) = w(z_j)$.*

Then there exists $x \in K$ with

$$v(x - x_i) > v(z_i) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Theorem 1.7. *Let $S_1, \dots, S_n \subseteq S_{\text{val}}(K)$, not necessarily disjoint.*

- (U) *Assume there exists a monic polynomial $f \in K[X]$ such that for every $v \in S_1 \cup \dots \cup S_n$, f has coefficients in the valuation ring of v and the reduction of f has no zero in the residue field of v .*
- (T) *Assume that each S_i is closed in the Hochster dual of the Zariski topology on $S_{\text{val}}(K)$.*

⁶See [Pre85, p. 354], also reproven in [FHV94, Corollary 1.3].

⁷The PRC property can indeed be seen as a very strong independence assumption, since it implies in particular that distinct orderings on K induce distinct topologies, cf. [Pre85, p. 353].

⁸See discussion in Section 5.7 how this follows from the results in [Ers01].

⁹We use ‘compact’ to mean what other sources call ‘quasi-compact’, i.e. there is no implication of being a Hausdorff space.

(I) Assume that for any i, j and $w \in S_i \cap S_j$, we have $w(x_i - x_j) \geq w(z_i) = w(z_j)$.

Then there exists $x \in K$ with

$$v(x - x_i) \geq v(z_i) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Theorem 1.8. Let $S_1, \dots, S_n \subseteq S_{\text{ord}}(K)$ pairwise disjoint.

(T) Assume that each S_i is compact in the Harrison topology on $S_{\text{ord}}(K)$.

(I) Assume that if w is a valuation on K whose valuation ring \mathcal{O}_w is convex both with respect to an ordering in S_i and with respect to an ordering in S_j , $i \neq j$, then these two orderings induce distinct orderings on the residue field of w , and $w(x_i - x_j) \geq w(z_i) = w(z_j)$.

Then there exists $x \in K$ with

$$(x - x_i)^2 < z_i^2 \text{ for all orderings in } S_i, \text{ for } i = 1, \dots, n.$$

Note that condition 1.8(I) is in particular always satisfied when distinct orderings on K induce distinct topologies, and so this in particular generalizes Theorem 1.4. In Remark 4.7, we explain how to deduce Theorems 1.6, 1.7 and 1.8 from Theorem 4.6, which is proven in sections 2–4.

In Section 5 we deduce from Theorem 4.6 some related results, namely value approximation theorems and residue approximation theorems. In the literature such results are often needed on the way to prove an approximation theorem, but we deduce them as corollaries of the main theorem. We also discuss special cases and applications to p -valuations and the connection to the strong approximation theorem in global fields.

In Section 6 we extend Theorem 4.6 even further by allowing finitely many exceptional valuations or (possibly complex) absolute values for which condition (U) might in general not be satisfiable, thus obtaining a result that in addition contains Theorem 1.1 and Theorem 1.3.

Finally, in Section 7 we use our approximation theorems to discuss a related problem, namely approximation of the values of rational functions. The results of Section 7 are crucial ingredients of our forthcoming paper [ADF18b], and the results of Section 4 will be used in [ADF18a].

2. THE SPACE OF LOCALITIES

We start by setting up the language unifying valuations and orderings in which we will phrase and prove the main theorem. For basics on valuations and orderings we refer to [EP05] and [Efr06].

Let K be a field. We denote by $\mathcal{B}(K)$ the set of subsets $\mathcal{O} \subseteq K$ that satisfy $\mathcal{O} \cdot \mathcal{O} \subseteq \mathcal{O}$ and $\mathcal{O} \cup \mathcal{O}^{-1} = K$ (where for $X \subseteq K$ we write $X^{-1} := \{x^{-1} : 0 \neq x \in X\}$).

Example 2.1. (1) $K \in \mathcal{B}(K)$

(2) If $|\cdot|$ is an absolute value on K , then $\{x \in K : |x| \leq 1\} \in \mathcal{B}(K)$.

(3) If v is a valuation on K with valuation ring \mathcal{O}_v , then $\mathcal{O}_v \in \mathcal{B}(K)$.

(4) If \leq is an ordering on K , then $\{x \in K : -1 \leq x \leq 1\} \in \mathcal{B}(K)$.

For $\mathcal{O} \in \mathcal{B}(K)$ we write $\mathcal{O}^\times = \mathcal{O} \cap \mathcal{O}^{-1}$ and $\mathfrak{m}_{\mathcal{O}} = \mathcal{O} \setminus \mathcal{O}^\times$. The following basic properties are easily checked: $1 \in \mathcal{O}$, $0 \in \mathfrak{m}_{\mathcal{O}}$, $\mathcal{O}\mathcal{O} = \mathcal{O}$, $\mathfrak{m}_{\mathcal{O}}\mathcal{O} = \mathfrak{m}_{\mathcal{O}}$. The collection of sets $x\mathcal{O}$, for $x \in K^\times$, is totally ordered by inclusion.

We equip $\mathcal{B}(K)$ with the Zariski topology \mathcal{T}_{Zar} with subbasis

$$\{\mathcal{O} \in \mathcal{B}(K) : x \in \mathcal{O}\}, x \in K.$$

The corresponding constructible topology \mathcal{T}_{con} by definition has subbasis

$$\{\mathcal{O} \in \mathcal{B}(K) : x \in \mathcal{O}\}, \{\mathcal{O} \in \mathcal{B}(K) : x \in \mathfrak{m}_{\mathcal{O}}\}, x \in K.$$

With the constructible topology, $\mathcal{B}(K)$ is a closed subspace of the Stone space 2^K , so it is compact and T_2 . Noting also that the Zariski topology is T_0 , it follows from [Hoc69, Proposition 7] that the Zariski topology is spectral, and it follows from [Hoc69, Proposition 8] that

the ‘‘Hochster dual’’ $\mathcal{T}_{\text{Zar}^*}$ of the Zariski topology with subbasis¹⁰

$$\{\mathcal{O} \in \mathcal{B}(K) : x \in \mathfrak{m}_{\mathcal{O}}, x \in K,\}$$

is spectral, see also [DST19, §1.4].

Remark 2.2. Consider the following properties of a subset $X \subseteq \mathcal{B}(K)$:

- (1) X is closed in \mathcal{T}_{Zar} or in $\mathcal{T}_{\text{Zar}^*}$
- (2) X is closed in \mathcal{T}_{con}
- (3) X is compact in \mathcal{T}_{con}
- (4) X is compact in \mathcal{T}_{Zar} and in $\mathcal{T}_{\text{Zar}^*}$

Then

$$(1) \Rightarrow (2) \Leftrightarrow (3) \Rightarrow (4).$$

We denote by $S_{\text{val}}(K)$ the space of equivalence classes of valuations on K , by $S_{\text{ord}}(K)$ the space of orderings on K , and by $S_{\text{abs}}(K)$ the space of equivalence classes of absolute values of K . We write

$$S(K) = S_{\text{val}}(K) \cup S_{\text{ord}}(K) \cup S_{\text{abs}}(K)$$

for the union of these three spaces, but identify rank-1 valuations and archimedean orderings with their associated absolute values. We call elements of $S(K)$ *localities*¹¹ of K and denote them by letters like v . In the case $v \in S_{\text{val}}(K)$, we denote by v also a fixed valuation in that equivalence class, in the case $v \in S_{\text{ord}}(K)$ we write \leq_v for the corresponding order relation on K , and in the case $v \in S_{\text{abs}}(K)$ we write $|\cdot|_v$ for a fixed absolute value in that equivalence class. As usual, we call an absolute value $v \in S_{\text{abs}}(K)$ *complex* if $|\cdot|_v$ is neither ultrametric nor induced by an archimedean ordering, i.e. if $v \in S_{\text{abs}}(K) \setminus (S_{\text{val}}(K) \cup S_{\text{ord}}(K))$.

For $v \in S(K)$ we write

$$\mathcal{O}_v = \begin{cases} \{x \in K : v(x) \geq 0\}, & v \in S_{\text{val}}(K), \\ \{x \in K : -1 \leq_v x \leq_v 1\}, & v \in S_{\text{ord}}(K), \\ \{x \in K : |x|_v \leq 1\}, & v \in S_{\text{abs}}(K) \end{cases}$$

and

$$\mathfrak{m}_v = \begin{cases} \{x \in K : v(x) > 0\}, & v \in S_{\text{val}}(K), \\ \{x \in K : -1 <_v x <_v 1\}, & v \in S_{\text{ord}}(K), \\ \{x \in K : |x|_v < 1\}, & v \in S_{\text{abs}}(K). \end{cases}$$

Note that these definitions agree if $v \in S_{\text{abs}}(K) \cap S_{\text{val}}(K)$ or $v \in S_{\text{abs}}(K) \cap S_{\text{ord}}(K)$. The map $v \mapsto \mathcal{O}_v$ gives an embedding of $S(K)$ into $\mathcal{B}(K)$, and we identify $S(K)$ with its image. Under this identification, the trivial valuation v_{trivial} is identified with $K \in \mathcal{B}(K)$, and for any locality v we have $\mathfrak{m}_v = \mathfrak{m}_{\mathcal{O}_v}$.

Remark 2.3. On $S_{\text{val}}(K)$, the Zariski topology induces the usual Zariski topology with subbasis

$$\{v \in S_{\text{val}}(K) : v(x) \geq 0\}, x \in K,$$

and the constructible topology induces the usual constructible (or patch) topology with subbasis

$$\{v \in S_{\text{val}}(K) : v(x) \geq 0\}, \{v \in S_{\text{val}}(K) : v(x) > 0\}, x \in K.$$

Both the Zariski topology and the constructible topology induce on $S_{\text{ord}}(K)$ the usual Harrison-topology (see for example [Lam05, VIII,§6]) with subbasis

$$\{v \in S_{\text{ord}}(K) : x \geq_v 0\}, x \in K,$$

¹⁰By definition, $\mathcal{T}_{\text{Zar}^*}$ has as a basis for the closed sets the compact open sets in \mathcal{T}_{Zar} , i.e. the finite unions of finite intersections of sets of the form $\{\mathcal{O} \in \mathcal{B}(K) : x \in \mathcal{O}\}$, for $x \in K$. It follows that $\{\mathcal{O} \in \mathcal{B}(K) : x \in \mathcal{O}\}$, for $x \in K$, is a subbasis for the closed sets of $\mathcal{T}_{\text{Zar}^*}$. This is equivalent to the description given above.

¹¹We deviate slightly from the terminology of [Efr06, Chapter 7], where only elements of $S_{\text{val}}(K) \cup S_{\text{ord}}(K)$ are called localities.

since for example $x \mapsto \frac{1-x}{1+x}$ exchanges the intervals $[-1, 1]$ and $[0, \infty)$. In particular, $S_{\text{val}}(K)$ and $S_{\text{ord}}(K)$ are compact and therefore closed in $S(K)$ in the constructible topology, while $S_{\text{abs}}(K)$ is in general not.

Example 2.4. For $\pi \in K^\times$ and $e \in \mathbb{N}$ we denote by $S_\pi^e(K) \subseteq S_{\text{val}}(K)$ the set of valuations v on K with discrete value group Γ_v (say \mathbb{Z} is a convex subgroup of Γ_v) and $0 < v(\pi) \leq e$. Then the three topologies \mathcal{T}_{Zar} , $\mathcal{T}_{\text{Zar}^*}$, and \mathcal{T}_{con} induce the same topology on $S_\pi^e(K)$. To see this, note that $v(x) \geq 0 \Leftrightarrow v(x^e \pi) > 0$, and $v(x) > 0 \Leftrightarrow v(x^e \pi^{-1}) \geq 0$, for all $v \in S_\pi^e(K)$ and all $x \in K^\times$. This applies in particular to sets $S \subseteq S_\pi^1(K)$ of valuations with a common uniformizer.

Moreover, $S_\pi^e(K)$ is \mathcal{T}_{con} -closed in $S_{\text{val}}(K)$, since $v \in S_{\text{val}}(K)$ is in $S_\pi^e(K)$ if and only if $v(\pi) > 0$ and for all $x \in K^\times$ we have either $v(x^{-1}) \geq 0$ or $v(x^e \pi^{-1}) \geq 0$. Since $S_{\text{val}}(K)$ is compact in the constructible topology, it follows that $S_\pi^e(K)$ is also compact.

Any locality $v \in S(K)$ induces on K a field topology, which we call the v -topology, defined by taking $\{z\mathfrak{m}_v : z \in K^\times\}$ as a basis for the filter of neighbourhoods of 0. The sets $B_v(x, z) := x + z\mathfrak{m}_v$ with $x \in K$, $z \in K^\times$ form a basis for this topology.

If $\mathcal{O}_v \subseteq \mathcal{O}_w$, then v is a *refinement* of w and w is a *coarsening* of v . This defines a partial order on $S(K)$, and v and w are *incomparable* if they are incomparable in this partial order.

Remark 2.5. If v, w are valuations then this terminology is standard; whereas if v is an ordering, then it has no proper refinement, and a valuation w is a coarsening of v if and only if \mathcal{O}_w is convex with respect to \leq_v . If $v \in S_{\text{abs}}(K) \setminus S_{\text{val}}(K)$, i.e. v is an archimedean absolute value (real or complex), then v has no proper refinement, and its only proper coarsening is the trivial valuation, so in particular v is incomparable to any other nontrivial element of $S(K)$.

Remark 2.6. Note that by definition v refines w if and only if v is in the Zariski closure of $\{w\}$, which is the case if and only if w is in the $\mathcal{T}_{\text{Zar}^*}$ -closure of $\{v\}$. In particular, a subspace of $S(K)$ satisfies the T_1 separation axiom with respect to either \mathcal{T}_{Zar} or $\mathcal{T}_{\text{Zar}^*}$ if and only if its elements are pairwise incomparable.

For every v and w in $S(K)$ there exists a finest common coarsening $v \vee w$ in $S(K)$. If w is a valuation coarsening v , then v induces a locality $\bar{v} \in S(Kw)$, where Kw denotes the residue field of w . Moreover, either v and \bar{v} are both valuations, or both orderings, or both complex absolute values (and w is trivial in this case).

Two localities $v, w \in S(K)$ are called *independent* if they induce distinct topologies on K . If both v and w are nontrivial, then this is the case if and only if $v \vee w$ is the trivial valuation. It follows from the fact that the set of coarsenings of a given locality is totally ordered that dependency is an equivalence relation on $S(K)$, and for any finite number of pairwise dependent nontrivial localities there is another nontrivial locality which is a coarsening of all of them.

We call v and w *strongly incomparable* if v and w are incomparable and induce distinct (and then automatically independent) localities on the residue field of their finest common coarsening $v \vee w$. Two sets $S_1, S_2 \subseteq S(K)$ are called *incomparable* (respectively *strongly incomparable*, or *independent*) if any element of S_1 is incomparable (respectively strongly incomparable or independent) to/from any element of S_2 .

Remark 2.7. Note that if two valuations are incomparable then they are automatically strongly incomparable, while the failure of strong incomparability for orderings is described by the Baer–Krull theorem [EP05, Theorem 2.2.5].

We have the following approximation theorem for pairwise independent localities, generalizing Theorem 1.2.

Theorem 2.8. *Let $v_1, \dots, v_n \in S(K)$ be all nontrivial, let $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$.*

(I) *Assume that v_1, \dots, v_n are pairwise independent.*

Then there exists $x \in K$ with

$$x - x_i \in z_i \mathfrak{m}_{v_i} \text{ for } i = 1, \dots, n.$$

Proof. If all v_i are in $S_{\text{val}}(K) \cup S_{\text{abs}}(K)$, then this is [PZ87, Corollary 4.2]. For any non-archimedean ordering v_i , consider its finest proper coarsening \tilde{v}_i , a nontrivial valuation. Since $z_i \mathfrak{m}_{\tilde{v}_i} \subseteq z_i \mathfrak{m}_{v_i}$, we can replace v_i by \tilde{v}_i and observe that the pairwise independence is preserved, thereby reducing to the situation without non-archimedean orderings. \square

The following is a generalization of a special case of Theorem 1.3:

Proposition 2.9. *Let $v_1, v_2 \in S(K)$ and let $z_1, z_2 \in K^\times$ such that for the finest common coarsening $w = v_1 \vee v_2$ we have $z_1 \mathcal{O}_w = z_2 \mathcal{O}_w$.*

- (1) *If v_1 and v_2 are strongly incomparable, then there exists $z \in K^\times$ with $z \in z_1 \mathfrak{m}_{v_1}$ and $z^{-1} \in z_2^{-1} \mathfrak{m}_{v_2}$.*
- (2) *If v_1 and v_2 are comparable or strongly incomparable (e.g. at least one of them is a valuation), then there exists $z \in K^\times$ with $z \in z_1 \mathcal{O}_{v_1}$ and $z^{-1} \in z_2^{-1} \mathcal{O}_{v_2}$.*

Proof. Assume without loss of generality that $z_1 = 1$. Note that then in particular $z_1, z_2 \in \mathcal{O}_w^\times$.

In case (1), w is a valuation and v_1 and v_2 induce independent $\bar{v}_1, \bar{v}_2 \in S(Kw)$. The nonempty set $\{\bar{z} \in Kw^\times : \bar{z}^{-1} \in \bar{z}_2^{-1} \mathfrak{m}_{\bar{v}_2}\}$ is open with respect to the \bar{v}_2 -topology on Kw and hence contains a ball $\bar{x} + \bar{z}'_2 \mathfrak{m}_{\bar{v}_2}$ with $\bar{x}, \bar{z}'_2 \in Kw^\times$. By Theorem 2.8 there exists $\bar{z} \in Kw$ with $\bar{z} \in \bar{z}_1 \mathfrak{m}_{\bar{v}_1}$ and $\bar{z} \in \bar{x} + \bar{z}'_2 \mathfrak{m}_{\bar{v}_2}$, so $\bar{z}^{-1} \in \bar{z}_2^{-1} \mathfrak{m}_{\bar{v}_2}$. If $z \in \mathcal{O}_w$ is any lift of \bar{z} , then $z \in z_1 \mathfrak{m}_{v_1}$ and $z^{-1} \in z_2^{-1} \mathfrak{m}_{v_2}$.

In case (2), if v_1 and v_2 are comparable, say $\mathcal{O}_{v_1} \subseteq \mathcal{O}_{v_2}$, then $w = v_2$ and thus $z_1 \mathcal{O}_{v_2} = z_2 \mathcal{O}_{v_2}$, so $z = z_1$ satisfies the claim in this case. Otherwise v_1 and v_2 are strongly incomparable and the claim then follows from the stronger claim (1). \square

Remark 2.10 (Shifting and scaling). The assumption that $z_1 = 1$ in the preceding proof is an example of two general simplification principles which will be used in our proofs in various places. Consider an approximation problem on two sets of localities S_1, S_2 , where we wish to find an element $x \in K$ such that $x - x_1 \in z_1 \mathcal{O}_v$ for all $v \in S_1$, and $x - x_2 \in z_2 \mathcal{O}_v$ for all $v \in S_2$, for some given elements $x_1, x_2 \in K$ and $z_1, z_2 \in K^\times$. If $x_1 = x_2$, then we have the trivial solution $x = x_1$, so we may assume $x_1 \neq x_2$. For any constant $c \in K$, if we have a solution x' for the modified approximation problem determined by $x'_1 = x_1 - c$ and $x'_2 = x_2 - c$, then $x = x' + c$ is a solution for the original approximation problem. This *shifting* of the problem can for instance be used to assume $x_1 = 0$ without loss of generality. Similarly, for any constant $d \in K^\times$, if there is a solution x' to the modified approximation problem given by $x'_1 = dx_1$, $x'_2 = dx_2$, $z'_1 = dz_1$, $z'_2 = dz_2$, then $x = d^{-1}x'$ is a solution to the original approximation problem. This *scaling*, together with previous shifting, can be used to reduce to the case $x_1 = 0$, $x_2 = 1$; alternatively (but usually not additionally), we can suppose that $z_1 = 1$, as in the proof of Proposition 2.9. The compatibility conditions (I) we impose in our theorems, see for instance 1.6(I), are all unaffected by scaling and shifting.

3. A UNIFORMITY CONDITION ON A SET OF LOCALITIES

We now describe the general (U) condition that we use in the main theorem and deduce some first consequences. Fix a field K and let $S \subseteq S(K)$. Consider the following assumption on S .

Assumption 3.1. There exists a polynomial $f \in K[X]$ of degree $d \geq 2$ with leading coefficient a_d and an element $t \in K^\times$ such that the following conditions are satisfied:

- (i) For any $v \in S$ and any $x \in \mathcal{O}_v$ we have $f(x) \in \mathcal{O}_v$.
- (ii) For any $v \in S$ and any $x \in \mathcal{O}_v$ we have $f(x) \notin t\mathfrak{m}_v$.
- (iii) For any $v \in S$ and any $x \notin \mathcal{O}_v$ we have $f(x) \in x^d(\mathcal{O}_v \setminus t\mathfrak{m}_v) \cap (a_d x^d + x^{d-1} \mathcal{O}_v)$.
- (iv) For any $v \in S$ we have $\mathcal{O}_v + \mathcal{O}_v \subseteq t^{-1} \mathcal{O}_v$.

Remark 3.2. Note that conditions (i, ii) as well as (iii) imply that $t \in \mathcal{O}_v$, and therefore condition (iv) is trivial if v is a valuation. Moreover, conditions (ii, iii) give that f has no zero in K . If v is a complex absolute value, then K is dense in its completion $\hat{K} \cong \mathbb{C}$; since f has a zero in \hat{K} , conditions (ii, iii) will never both be satisfied in that case. We will therefore mostly ignore complex absolute values until Section 6.

Condition (iii) implies that $a_d \in \mathcal{O}_v$ for all $v \in S$: If v is a valuation and $a_d \notin \mathcal{O}_v$, then for any $x \notin \mathcal{O}_v$ we have $x^d \mathcal{O}_v \cap (a_d x^d + x^{d-1} \mathcal{O}_v) = \emptyset$, since a member of the intersection would have to have valuation both equal to $v(a_d x^d)$ and at least $v(x^d)$. If v is an ordering (and very similar if v is a complex absolute value) and $a_d \notin \mathcal{O}_v$, then for $x >_v \max(1, (|a_d| - 1)^{-1})$ we have $x^d \mathcal{O}_v + x^{d-1} \mathcal{O}_v \subseteq x^d(1 + x^{-1})\mathcal{O}_v$, and all elements of this set are of smaller absolute value than $a_d x^d$, so again $x^d \mathcal{O}_v \cap (a_d x^d + x^{d-1} \mathcal{O}_v) = \emptyset$.

Remark 3.3. Note that 3.1 remains true when S is enlarged by adding the coarsenings of all localities in S ; we may thus assume that S is closed under coarsenings. The assumption is also true for any $S' \subseteq S$.

Remark 3.4. Note that in the situation of Assumption 3.1, if $t \in \mathcal{O}_v^\times$ for some $v \in S$, then conditions (i,ii) imply that v is necessarily a valuation. In particular, if $t = 1$, then S consists only of valuations.

Example 3.5. Assumption 3.1 is satisfied for the following sets $S \subseteq S(K)$:

- (1) If $S \subseteq S_{\text{val}}(K)$ and there exists a monic $f \in K[X]$ such that for all valuations $v \in S$ the coefficients of f are in the valuation ring of v and the reduction of f does not have a zero in the residue field of v , then the assumption is satisfied with $t = 1$.
- (2) If $S \subseteq S_\pi^1(K)$ is a set of valuations with a common uniformizer $\pi \in K^\times$ (cf. Example 2.4), then the assumption is satisfied with $f = X^2 - \pi$ and $t = \pi$. Slightly more general, assume that $S \subseteq S_\pi^e(K)$, for some $\pi \in K^\times$ and $e > 0$ (also cf. Example 2.4). Then the hypothesis is satisfied with $f = X^{e+1} - \pi$ and $t = \pi$.
- (3) If $S \subseteq S_{\text{ord}}(K)$ then the hypothesis is satisfied with $f = \frac{1}{2}(X^2 + 1)$ and $t = \frac{1}{2}$.

Example 3.6. Assume that K is of characteristic zero. For a prime number p and $e \in \mathbb{N}$, we note that $S_p^e(K)$ contains all p -valuations in the sense of [PR84] with p -ramification index at most e (and arbitrary residue degree). Then for every finite set of prime numbers \mathcal{P} and $e \in \mathbb{N}$, the space

$$S = S_{\text{ord}}(K) \cup \bigcup_{p \in \mathcal{P}} S_p^e(K)$$

satisfies Assumption 3.1: With $q = \prod_{p \in \mathcal{P}} p$ we can take $f = \frac{q+1}{2q+1}(X^{2e} + \frac{q}{q+1})$ and $t = \frac{q}{2q+1}$. In particular, in the language of [Feh13], if S_0 is a finite set of primes of a number field K_0 and K is an extension of K_0 , then $S_{S_0}^\tau(K)$ satisfies Assumption 3.1 for any type $\tau \in \mathbb{N}^2$.

Example 3.7. As a different criterion, let $S \subseteq S_{\text{ord}}(K)$, $f = \sum_{i=0}^d a_i X^i \in \mathbb{Q}[X]$ and $t \in \mathbb{Q}^\times$. If

$$\sum_{i=0}^d |a_i| \leq 1, \quad |t| + \sum_{i=0}^{d-1} |a_i| \leq |a_d|, \quad \text{and} \quad |t| \leq \frac{1}{2},$$

then Conditions 3.1(i,iii,iv) are satisfied. If additionally $|f(x)| \geq |t|$ for every $x \in \mathbb{R}$, then the same holds for every real-closed field and therefore for every ordered field, so Condition 3.1(ii) is also satisfied.

Example 3.8. Let $g \in \mathbb{Z}[X]$ be a monic polynomial with no real zero, so in particular g as a function $\mathbb{R} \rightarrow \mathbb{R}$ is bounded away from zero. Write $g = \sum_{i=0}^d a_i X^i$. For a small rational number $b > 0$, consider the polynomial

$$f = b^{d+1} g(b^{-1} X) = b \cdot \sum_{i=0}^d a_i b^{d-i} X^i \in \mathbb{Q}[X]$$

and $t = b^{d+2}$. For sufficiently small b , this clearly satisfies all the conditions from Example 3.7. We may in fact choose $b = 1/l$ for some large prime number l .

Now let K be any field of characteristic not l , and consider f and t as above. In this situation Assumption 3.1 is satisfied with $S = S_{\text{ord}}(K) \cup S'$, where $S' \subseteq S_{\text{val}}(K)$ is the set of valuations on K with residue characteristic not l in whose residue field the reduction of g does not have

a zero: For any valuation $v \in S'$, the polynomial f has all coefficients in \mathcal{O}_v , its reduction has no zero over the residue field since the reduction of g does not, and $t \in \mathcal{O}_v^\times$. For orderings $v \in S_{\text{ord}}(K)$, we have forced the assumption to be satisfied by construction.

Note that by the Chebotarev Density Theorem (see [Neu99, VII, (13.4)]), the set S' will include all valuations on K with residue field \mathbb{F}_p for some positive density of prime numbers p . By choosing g suitably, e.g. as the minimal polynomial of an integral primitive element of a totally imaginary Galois extension L/\mathbb{Q} , one can in fact achieve densities arbitrarily close to 1, because the reduction of the polynomial g has no root in \mathbb{F}_p for all prime numbers p such that p is neither completely split nor ramified in L/\mathbb{Q} , cf. [Neu99, I, (8.3)].

Proposition 3.9. *Under Assumption 3.1(i, ii, iii), there exists a function $\phi : K^\times \times K^\times \longrightarrow K^\times$ such that for $x, y \in K^\times$ and $v \in S$ the following hold:*

- (1) *If $x \in t\mathcal{O}_v$ and $y \in t\mathcal{O}_v$, then $\phi(x, y) \in x\mathcal{O}_v \cup y\mathcal{O}_v$.*
- (2) *If $x^{-1} \in \mathcal{O}_v$ or $y^{-1} \in \mathcal{O}_v$, then $\phi(x, y)^{-1} \in x^{-1}\mathcal{O}_v \cap y^{-1}\mathcal{O}_v$.*

Proof. Let ϕ be the homogenisation of ft^{-1} , so if f is of degree $d \geq 2$ then ϕ is given by

$$\phi(X, Y) = f(XY^{-1})Y^dt^{-1} \in K[X, Y].$$

Let $v \in S$ and write $\mathcal{O} = \mathcal{O}_v$, $\mathfrak{m} = \mathfrak{m}_v$.

To prove (1), let $x, y \in t\mathcal{O}$. Then $x^dt^{-1} \in x\mathcal{O}$ and $y^dt^{-1} \in y\mathcal{O}$. If $xy^{-1} \in \mathcal{O}$, then $f(xy^{-1}) \in \mathcal{O}$, by 3.1(i), hence

$$\phi(x, y) = f(xy^{-1})y^dt^{-1} \in y\mathcal{O}.$$

If instead $xy^{-1} \notin \mathcal{O}$, then $f(xy^{-1}) \in (xy^{-1})^d\mathcal{O}$ by 3.1(iii), hence

$$\phi(x, y) = f(xy^{-1})y^dt^{-1} \in (xy^{-1})^dy^dt^{-1}\mathcal{O} = x^dt^{-1}\mathcal{O} \subseteq x\mathcal{O},$$

as required.

To prove (2), our assumption is that either $x^{-1} \in \mathcal{O}$ or $y^{-1} \in \mathcal{O}$. If $xy^{-1} \in \mathcal{O}$, then $y^{-1}\mathcal{O} \subseteq x^{-1}\mathcal{O}$, so the assumption implies that $y^{-1} \in \mathcal{O}$. From 3.1(ii) we have $f(xy^{-1}) \notin \mathfrak{m}$, so $f(xy^{-1})^{-1} \in t^{-1}\mathcal{O}$. Thus

$$\phi(x, y)^{-1} = f(xy^{-1})^{-1}y^{-dt} \in y^{-d}\mathcal{O} \subseteq y^{-1}\mathcal{O} = x^{-1}\mathcal{O} \cap y^{-1}\mathcal{O}.$$

On the other hand, if $xy^{-1} \notin \mathcal{O}$, then $x^{-1}y \in \mathcal{O}$ and $x^{-1}\mathcal{O} \subseteq y^{-1}\mathcal{O}$, and the assumption implies that $x^{-1} \in \mathcal{O}$. By 3.1(iii), $f(xy^{-1}) \notin (xy^{-1})^d\mathfrak{m}$, so $f(xy^{-1})^{-1} \in (x^{-1}y)^dt^{-1}\mathcal{O}$. Thus

$$\phi(x, y)^{-1} = f(xy^{-1})^{-1}y^{-dt} \in (x^{-1}y)^dt^{-1}y^{-dt}\mathcal{O} = x^{-d}\mathcal{O} \subseteq x^{-1}\mathcal{O} = x^{-1}\mathcal{O} \cap y^{-1}\mathcal{O}. \quad \square$$

Corollary 3.10. *Under Assumption 3.1(i, ii, iii), and with ϕ as in Proposition 3.9, for $x_1, \dots, x_n \in K^\times$, the nonzero element*

$$\phi(x_1, \dots, x_n) := \begin{cases} x_1, & n = 1 \\ \phi(x_1, \phi(x_2, \dots)), & n > 1 \end{cases}$$

satisfies for every $v \in S$:

- (1) *If $x_i \in t\mathcal{O}_v$ for all i , then $\phi(x_1, \dots, x_n) \in \bigcup_{i=1}^n x_i\mathcal{O}_v$.*
- (2) *If $x_i^{-1} \in \mathcal{O}_v$ for some i , then $\phi(x_1, \dots, x_n)^{-1} \in \bigcap_{i=1}^n x_i^{-1}\mathcal{O}_v$.*

Proof. For (1), induction using Proposition 3.9(1) firstly shows that for any $i \geq 1$, the element $\phi(x_i, \dots, x_n)$ is in $t\mathcal{O}_v$. Secondly, using Proposition 3.9(1) again, we obtain

$$\phi(x_1, \dots, x_n) = \phi(x_1, \phi(x_2, \dots, x_n)) \in x_1\mathcal{O}_v \cup \phi(x_2, \dots, x_n)\mathcal{O}_v,$$

so using induction once more gives $\phi(x_1, \dots, x_n) \in \bigcup_i x_i\mathcal{O}_v$ as desired.

For (2) note that it is sufficient to prove that $\phi(x_1, \dots, x_n)^{-1} \in x_i^{-1}\mathcal{O}_v$ for those i with $x_i^{-1} \in \mathcal{O}_v$, since for $x_i^{-1} \in \mathcal{O}_v$ and $x_{i'}^{-1} \notin \mathcal{O}_v$ we have $x_i^{-1}\mathcal{O}_v \subseteq x_{i'}^{-1}\mathcal{O}_v$. For any such i , we have $\phi(x_i, \dots, x_n)^{-1} \in x_i^{-1}\mathcal{O}_v$ by Proposition 3.9(2), and then induction shows $\phi(x_j, \dots, x_n)^{-1} \in x_i^{-1}\mathcal{O}_v$ for all $j \leq i$, so we obtain the claim $\phi(x_1, \dots, x_n)^{-1} \in x_i^{-1}\mathcal{O}_v$. \square

Remark 3.11. Only for use in Section 6 we note that the proof of part (1) of Proposition 3.9, and consequently also of part (1) of Corollary 3.10, requires only 3.1(i) and 3.1(iii) to hold.

Remark 3.12. A function ϕ satisfying the conditions from Proposition 3.9 can also be obtained using assumptions different from our Assumption 3.1. For instance, if one works with a set S consisting exclusively of valuations such that the holomorphy ring $R = \bigcap_{v \in S} \mathcal{O}_v$ is a Bézout ring with quotient field K , then for any $x, y \in K^\times$ we can find $z \in K^\times$ with $v(z) = \min(v(x), v(y))$ for all $v \in S$: this is exactly the statement that the fractional ideal $xR + yR$ is principal with generator z . This z will satisfy the properties desired from $\phi(x, y)$ in Proposition 3.9.

As a slightly more general condition, it is sufficient for R to be a Prüfer ring such that there exists a natural number n for which the n -th power of any two-generated fractional ideal of R is principal: In this situation, for any $x, y \in K^\times$, we can find a generator z of the fractional ideal $(xR + yR)^n$, and this generator z will satisfy $v(z) = n \min(v(x), v(y))$ for all $v \in S$, which is as desired for $\phi(x, y)$. This latter condition on R is established in [Roq73, Theorem 1] in the situation of Example 3.5(1), using effectively our construction of the function ϕ .

Ring-theoretic conditions on the holomorphy ring are in the style of the (U) conditions of results of Ershov, see Section 5.7. Our Assumption 3.1 is more ad hoc, but has the advantage that it can also apply to orderings.

4. THE APPROXIMATION THEOREM

We want to prove two approximation theorems: one with a conclusion of the type $x - x_i \in z_i \mathfrak{m}$ and one with the weaker conclusion $x - x_i \in z_i \mathcal{O}$. Neither of these theorems will easily imply the other, but the proofs are very similar, so we state and prove our lemmas simultaneously in the two situations, called “Situation \mathfrak{m} ” and “Situation \mathcal{O} ”.

Let $S \subseteq S(K)$. For $v \in S(K)$, we write $\mathfrak{B}_v = \mathfrak{m}_v$ in Situation \mathfrak{m} , and $\mathfrak{B}_v = \mathcal{O}_v$ in Situation \mathcal{O} . Moreover, in Situation \mathfrak{m} , we endow $S(K)$ with the topology $\mathcal{T}_{\text{Zar}^*}$, in Situation \mathcal{O} with the Zariski topology \mathcal{T}_{Zar} . In this way, in either case sets of the form $\{v \in S(K) : x \in \mathfrak{B}_v\}$, for $x \in K$, are open. Also, for the rest of this section we assume that S satisfies Assumption 3.1. Indeed, we fix a polynomial $f \in K[X]$ of degree $d \geq 2$ and an element $t \in K^\times$, with respect to which S satisfies this assumption. We call two localities v, w *t-incomparable* if $t \in \mathcal{O}_{v \vee w}^\times$ and additionally v, w are strongly incomparable if they are both orderings. Note that any valuation is 1-incomparable to itself. We call two sets S_1, S_2 of localities *t-incomparable* if any element of S_1 is *t-incomparable* to any element of S_2 .

Example 4.1. Let $t = \pi \in K$ and $e \in \mathbb{N}$. Valuations $v, w \in S_\pi^e(K)$ (cf. Example 2.4) are *t-incomparable* if and only if they are incomparable, which is the case if and only if they are distinct.

We now start the proof of the approximation theorem by constructing elements that are ‘uniformly small’ resp. ‘uniformly big’.

Lemma 4.2. *Let $S_1, S_2 \subseteq S$ be nonempty and compact in the given topology. Assume they are t-incomparable, and in Situation \mathfrak{m} furthermore incomparable. Let $z, z' \in K^\times$ such that $w(z) \leq 0$ and $w(z') \leq 0$ whenever w is a valuation on K which has a refinement in S_1 and a refinement in S_2 . Then there exists $b \in K^\times$ such that $b \in t\mathfrak{B}_v \cap tz'\mathfrak{B}_v$ for all $v \in S_1$, and $b^{-1} \in t\mathfrak{B}_v \cap tz'\mathfrak{B}_v$ for all $v \in S_2$.*

Proof. For fixed $v \in S_1$ and $v' \in S_2$, denote $w = v \vee v'$. This is a valuation by the assumption of *t-incomparability*, and so $w(z), w(z') \leq 0$, and $w(t) = 0$.

With the aim of applying Proposition 2.9, we define two elements $z_v, z'_{v'}$ as modifications of z, z' , as follows. If $z \in \mathcal{O}_v$ let $z_v = tz$, otherwise let $z_v = t$. If $z' \in \mathcal{O}_{v'}$ let $z'_{v'} = (tz')^{-1}$, otherwise let $z'_{v'} = t^{-1}$. Note that $w(z_v) = w(z'_{v'}) = 0$. Therefore $z_v \mathcal{O}_w = \mathcal{O}_w = z'_{v'} \mathcal{O}_w$, so the assumption of Proposition 2.9 is satisfied.

In Situation \mathfrak{m} we have that v and v' are strongly incomparable and can apply Proposition 2.9(1) to get $y \in K^\times$ with $y \in z_v \mathfrak{m}_v = z_v \mathfrak{B}_v$ and $y^{-1} \in z'_{v'} \mathfrak{m}_{v'} = z'_{v'} \mathfrak{B}_{v'}$; In Situation \mathcal{O} we can

apply Proposition 2.9(2) to get $y \in K^\times$ with $y \in z_v \mathcal{O}_v = z_v \mathfrak{B}_v$ and $y^{-1} \in z_{v'}^{-1} \mathcal{O}_{v'} = z_{v'}^{-1} \mathfrak{B}_{v'}$. Thus, setting $b_{vv'} = y$, we get in both situations that

- (1) $b_{vv'} \in z_v \mathfrak{B}_v = t \mathfrak{B}_v \cap tz \mathfrak{B}_v$ and
- (2) $b_{vv'}^{-1} \in z_{v'}^{-1} \mathfrak{B}_{v'} = t \mathfrak{B}_{v'} \cap tz' \mathfrak{B}_{v'}$.

For every $x \in K^\times$, the set

$$U_x^{(2)} = \{u \in S : x^{-1} \in t \mathfrak{B}_u \cap tz' \mathfrak{B}_u\} = \{u \in S : (xt)^{-1} \in \mathfrak{B}_u\} \cap \{u \in S : (xtz')^{-1} \in \mathfrak{B}_u\}$$

is open in the given topology. For each $v \in S_1$, the family $\{U_{b_{vv'}}^{(2)} : v' \in S_2\}$ is an open covering of S_2 , by (2). By compactness of S_2 , there are finitely many $v'_1, \dots, v'_m \in S_2$ such that $S_2 \subseteq \bigcup_{i=1}^m U_{b_{vv'_i}}^{(2)}$. Let $b_v = \phi(b_{vv'_1}, \dots, b_{vv'_m})$. For each i , by (1), we have $b_{vv'_i} \in t \mathfrak{B}_v \cap tz \mathfrak{B}_v \subseteq t \mathcal{O}_v$, and so Corollary 3.10(1) gives that

$$(1') \quad b_v \in \bigcup_{i=1}^m b_{vv'_i} \mathcal{O}_v \subseteq t \mathfrak{B}_v \cap tz \mathfrak{B}_v.$$

On the other hand, for each $v' \in S_2$, there is an i with $v' \in U_{b_{vv'_i}}^{(2)}$, i.e. $b_{vv'_i}^{-1} \in t \mathfrak{B}_{v'} \cap tz' \mathfrak{B}_{v'} \subseteq \mathcal{O}_{v'}$, and so Corollary 3.10(2) gives that

$$(2') \quad b_v^{-1} \in \bigcap_{i=1}^m b_{vv'_i}^{-1} \mathcal{O}_{v'} \subseteq t \mathfrak{B}_{v'} \cap tz' \mathfrak{B}_{v'}.$$

Similarly, for every $x \in K$, the set

$$U_x^{(1)} = \{u \in S : x \in t \mathfrak{B}_u \cap tz \mathfrak{B}_u\}$$

is open, and the family $\{U_{b_v}^{(1)} : v \in S_1\}$ is an open covering of S_1 , by (1'). By compactness of S_1 , there exist finitely many $v_1, \dots, v_k \in S_1$ such that $S_1 \subseteq \bigcup_{i=1}^k U_{b_{v_i}}^{(1)}$. Let $b = \phi(b_{v_1}^{-1}, \dots, b_{v_k}^{-1})^{-1}$.

For each $v \in S_1$, there is an i with $v \in U_{b_{v_i}}^{(1)}$, i.e. $(b_{v_i}^{-1})^{-1} = b_{v_i} \in t \mathfrak{B}_v \cap tz \mathfrak{B}_v \subseteq \mathcal{O}_v$ and so Corollary 3.10(2) gives that

$$(1'') \quad b \in \bigcap_{i=1}^k b_{v_i} \mathcal{O}_v \subseteq t \mathfrak{B}_v \cap tz \mathfrak{B}_v.$$

Finally, for each $v' \in S_2$ and each i , by (2'), we have that $b_{v_i}^{-1} \in t \mathfrak{B}_{v'} \cap tz' \mathfrak{B}_{v'} \subseteq t \mathcal{O}_{v'}$, and so Corollary 3.10(1) gives that

$$(2'') \quad b^{-1} \in \bigcup_{i=1}^k b_{v_i}^{-1} \mathcal{O}_{v'} \subseteq t \mathfrak{B}_{v'} \cap tz' \mathfrak{B}_{v'},$$

as required. \square

Lemma 4.3. *Let f, t, d, a_d as in Assumption 3.1. Let $v \in S$, $z \in K^\times$ and $b \in K^\times$. Then the element $x = a_d b^{-d} f(b^{-1})^{-1}$ satisfies*

- (1) $x - 1 \in z \mathfrak{B}_v$ if $b \in t \mathfrak{B}_v \cap tz \mathfrak{B}_v$, and
- (2) $x \in z \mathfrak{B}_v$ if $b^{-1} \in t \mathfrak{B}_v \cap tz \mathfrak{B}_v$.

Proof. Recall that $t \in \mathcal{O}_v$ and $a_d \in \mathcal{O}_v$ (Remark 3.2).

Proof of (1): Assume that $b \in t \mathfrak{B}_v \cap tz \mathfrak{B}_v$. If $b \in \mathfrak{m}_v$ (which is the case if $t \in \mathfrak{m}_v$, $z \in \mathfrak{m}_v$, or $\mathfrak{B}_v = \mathfrak{m}_v$), then $b^d f(b^{-1}) \in (\mathcal{O}_v \setminus t \mathfrak{m}_v) \cap (a_d + b \mathcal{O}_v)$ by 3.1(iii), so in particular $a_d - b^d f(b^{-1}) \in b \mathcal{O}_v$ and $(b^d f(b^{-1}))^{-1} \in t^{-1} \mathcal{O}_v$. Therefore,

$$x - 1 = \frac{a_d - b^d f(b^{-1})}{b^d f(b^{-1})} \in b \mathcal{O}_v t^{-1} \mathcal{O}_v = b t^{-1} \mathcal{O}_v \subseteq z \mathfrak{B}_v$$

as desired. If $b \notin \mathfrak{m}_v$, then $\mathfrak{B}_v = \mathcal{O}_v$, $t \in \mathcal{O}_v^\times$ and $z^{-1} \in \mathcal{O}_v$. Combining $b \in t \mathfrak{B}_v = \mathcal{O}_v$ and $b^{-1} \in \mathcal{O}_v$ we then get $b \in \mathcal{O}_v^\times$. Therefore, $b^d f(b^{-1}) \in \mathcal{O}_v \setminus t \mathfrak{m}_v = \mathcal{O}_v^\times$ by 3.1(i,ii), and therefore $x \in \mathcal{O}_v$. Since $t \in \mathcal{O}_v^\times$ implies that v is a valuation (Remark 3.4), we conclude that

$$x - 1 \in \mathcal{O}_v \subseteq z \mathcal{O}_v = z \mathfrak{B}_v.$$

Proof of (2): Now assume that $b^{-1} \in t \mathfrak{B}_v \cap tz \mathfrak{B}_v$. We have $f(b^{-1}) \notin t \mathfrak{m}_v$ by 3.1(ii), so

$$x \in a_d b^{-d} t^{-1} \mathcal{O}_v \subseteq b^{-d+1} \mathfrak{B}_v \subseteq z \mathfrak{B}_v. \quad \square$$

Remark 4.4. Again only for use in Section 6 we point out that the proof of Lemma 4.3(1) in Situation **m** requires only 3.1(iii) to hold.

Lemma 4.5. *Let $S_1, \dots, S_n \subseteq S$ be nonempty and compact in the given topology. Assume they are pairwise t -incomparable, and in Situation \mathfrak{m} furthermore pairwise incomparable. Let $x', x'' \in K$, $z_1, \dots, z_n \in K^\times$ such that for any valuation w on K which has a refinement in S_i and a refinement in S_j , $i \neq j$, we have $w(z_i) = w(z_j)$, and furthermore for any valuation w with a refinement in S_1 and a refinement in S_i , $i \neq 1$, we have $w(x' - x'') \geq w(z_1)$. Then there exists $x \in K$ with $x - x' \in z_1 \mathfrak{B}_v$ for each $v \in S_1$ and $x - x'' \in z_i \mathfrak{B}_v$ for each $v \in S_i$, $i \neq 1$.*

Proof. If $x' = x''$, we may take $x = x'$, so assume this is not the case. By shifting and scaling as in Remark 2.10, we may assume that $x'' = 0$ and $x' = 1$. (Observe that all the hypotheses and the claim are invariant under shifting and scaling.) In this situation, the compatibility hypothesis implies that $w(z_1) = w(z_i) \leq 0$ for every w with a refinement in S_1 and a refinement in S_i , $i \neq 1$.

For any $i \neq 1$, Lemma 4.2 provides an element $b_i \in K^\times$ with $b_i \in t\mathfrak{B}_v \cap tz_1\mathfrak{B}_v$ for $v \in S_1$ and $b_i^{-1} \in t\mathfrak{B}_v \cap tz_i\mathfrak{B}_v$ for $v \in S_i$. Let $b = \phi(b_2, \dots, b_n)$. By Corollary 3.10(1,2), $b \in t\mathfrak{B}_v \cap tz_1\mathfrak{B}_v$ for $v \in S_1$, and $b^{-1} \in t\mathfrak{B}_v \cap tz_i\mathfrak{B}_v$ for $v \in S_i$, $i \neq 1$. Hence Lemma 4.3 finishes the proof. \square

Theorem 4.6. *Let $S_1, \dots, S_n \subseteq S(K)$, $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$.*

- (U) *Assume that Assumption 3.1 holds for $S = S_1 \cup \dots \cup S_n$ for some f, t .*
- (T) *Assume each S_i is compact in the given topology.*
- (I) *Assume that for any valuation w on K with a refinement in S_i and a refinement in S_j we have $w(x_i - x_j) \geq w(z_i) = w(z_j)$; assume further that the S_i are t -incomparable, and in Situation \mathfrak{m} furthermore incomparable.*

Then there exists $x \in K$ with

$$x - x_i \in z_i \mathfrak{B}_v \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Proof. We use induction on n . Without loss of generality assume that $S_i \neq \emptyset$ for all i .

For $n = 1$ we may take $x = x_1$.

For $n > 1$, we want to apply the induction hypothesis for the sets S_2, \dots, S_n , the elements x_2, \dots, x_n and tz_2, \dots, tz_n . Note that, by the assumption of pairwise t -incomparability, $w(tz_i) = w(z_i)$ for every valuation w with a refinement in S_i and a refinement in S_j , $j \neq i$, so the compatibility condition $w(x_i - x_j) \geq w(tz_i) = w(tz_j)$ is satisfied. The induction hypothesis thus gives an element $x'' \in K$ with $x'' - x_i \in z_i t\mathfrak{B}_v$ for all $v \in S_i$, $i \geq 2$. We now apply Lemma 4.5 to the sets S_1, \dots, S_n and the elements $x' := x_1$, x'' and $z_1, z_2 t, \dots, z_n t$. Note that if a valuation w has a refinement in S_1 and a refinement v in S_i , $i \neq 1$, then indeed

$$w(x' - x'') = w((x_1 - x_i) + (x_i - x'')) \geq w(z_1) = w(z_i),$$

as $x'' - x_i \in z_i t\mathfrak{B}_v \subseteq z_i \mathcal{O}_w$ implies that $w(x'' - x_i) \geq w(z_i)$. We thus obtain $x \in K$ with $x - x_1 \in z_1 \mathfrak{B}_v$ for all $v \in S_1$, and $x - x'' \in z_i t\mathfrak{B}_v$ for all $v \in S_i$, $i \neq 1$. Finally, for $v \in S_i$, $i \neq 1$, we get that

$$x - x_i = (x - x'') + (x'' - x_i) \in z_i t\mathfrak{B}_v + z_i t\mathfrak{B}_v \subseteq z_i t\mathfrak{B}_v (\mathcal{O}_v + \mathcal{O}_v) \subseteq z_i t\mathfrak{B}_v \cdot t^{-1} \mathcal{O}_v = z_i \mathfrak{B}_v$$

by 3.1(iv). This finishes the induction. \square

Remark 4.7. As promised, we now explain how to deduce Theorems 1.6, 1.7 and 1.8 from the introduction.

Theorem 1.6 is obtained by applying Theorem 4.6 in Situation \mathfrak{m} with $t = \pi$: 1.6(U) implies 4.6(U) (see Example 3.5(2)), and 1.6(T) together with 1.6(U) implies 4.6(T) (see Example 2.4). The S_i are pairwise t -incomparable and pairwise incomparable since pairwise disjoint (Example 4.1), so 1.6(I) implies 4.6(I).

Theorem 1.7 is obtained by applying Theorem 4.6 in Situation \mathcal{O} with $t = 1$: 1.7(U) implies 4.6(U) (see Example 3.5(1)), and 1.7(T) implies 4.6(T) (see Remark 2.2). In fact, since the S_i are closed in the Hochster dual of the Zariski topology, then the S_i are also closed under coarsenings, and so 1.7(I) implies 4.6(I).

Finally, Theorem 1.8 follows from Theorem 4.6 in Situation \mathfrak{m} using Example 3.5(3) and taking Remark 2.3 into consideration.

Remark 4.8. Note that the set of $v \in S(K)$ for which a given $x \in K$ satisfies an approximation condition $x - x_i \in z_i \mathfrak{B}_v$ is always open-closed in the constructible topology, in particular compact both in \mathcal{T}_{Zar} and in $\mathcal{T}_{\text{Zar}^*}$. This explains why condition 4.6(T) is natural. It is also clear that without this condition the theorem must fail.

Remark 4.9. It is obvious that assumption 4.6(I) cannot simply be dropped. It is also clear that if there exists x with $x - x_i \in z_i \mathcal{O}_v$ for all $v \in S_i$, $i = 1, \dots, n$, then any w as in 4.6(I) must satisfy $w(x_i - x_j) \geq \min\{w(z_i), w(z_j)\}$, but the condition $w(x_i - x_j) \geq w(z_i) = w(z_j)$, which also appears in Theorem 1.3, cannot be deduced and could possibly be relaxed. However, the following example shows that one cannot replace this compatibility condition by $w(x_i - x_j) \geq \min\{w(z_i), w(z_j)\}$ in Theorem 4.6:

If $K = \mathbb{Q}(T)$ with w the T -adic valuation, and v_1 and v_2 the composites of w with the 2- and 3-adic valuations, respectively, then there is no x with $v_1(x - (2T)^{-1}) \geq v_1(T^{-1})$ and $v_2(x) \geq 0$, since $v_1(x) = v_1((2T)^{-1})$ implies $w(x) = -1$, but $v_2(x) \geq 0$ would imply $w(x) \geq 0$. In this case we might have sought to apply Theorem 4.6 in Situation \mathcal{O} with $S_1 = \{v_1\}$, $S_2 = \{v_2\}$, $x_1 = (2T)^{-1}$, $x_2 = 0$, $z_1 = T^{-1}$, and $z_2 = 1$. Then assumption (U) is satisfied by 3.5(1), and $w(x_1 - x_2) = -1 = \min\{w(z_1), w(z_2)\}$.

Remark 4.10. We can conclude from Theorem 1.8 that every field K for which

- (*) any two orderings on K induce distinct orderings on the residue field of their finest common coarsening w

is an SAP-field in the sense of [Pre84, §6], i.e. for every two disjoint closed subsets S_1 and S_2 of $S_{\text{ord}}(K)$ there is some $a \in K$ with $a >_v 0$ for all $v \in S_1$ and $a <_v 0$ for all $v \in S_2$. This is in fact a special case of [Pre84, Theorem 9.1], which in particular states that K is SAP if and only if for every valuation w on K with formally real residue field either

- (i) the value group Γ_w is 2-divisible, or
- (ii) $|\Gamma_w/2\Gamma_w| = 2$ and the residue field Kw carries a unique ordering.

Due to the Baer-Krull theorem, our condition (*) is precisely that (i) holds always. We note that Theorem 1.8, and therefore also Theorem 4.6, is no longer true if we replace (*) by the weaker condition that K is an SAP-field, as the example $K = \mathbb{R}((T))$ with its two orderings \leq_{0^+} and \leq_{0^-} shows: The approximation problem with $S_1 = \{\leq_{0^+}\}$, $S_2 = \{\leq_{0^-}\}$, $x_1 = 0$, $x_2 = 1$, $z_1 = z_2 = \frac{1}{3}$ satisfies the compatibility condition $w(x_1 - x_2) \geq w(z_1) = w(z_2)$ where w is the T -adic valuation, but is not solvable.

5. APPLICATIONS AND COUNTEREXAMPLES

We start by deducing a few corollaries that resemble similar approximation theorems in the literature. We will phrase several of these corollaries for compact sets in $\mathcal{T}_{\text{Zar}^*}$, but recall that this property is satisfied for example by every closed set in the constructible topology (Remark 2.2).

5.1. Value approximation. The following ‘value approximation’ theorem is our version of Ribenboim’s [Rib57, Theorem 5], see also [Rib68, p. 135 Théorème 1] and [War89, Theorem 28.12], which in the case of independent valuations appears already in Krull’s seminal paper [Kru31, Satz 15]. In [Ers01, Proposition 2.6.6] a similar result with a different condition (U) is given, and condition (T) is replaced by compactness in the Zariski topology.

Corollary 5.1. *Let $S_1, \dots, S_n \subseteq S_{\text{val}}(K)$ and $z_1, \dots, z_n \in K^\times$.*

- (U) *Assume that Assumption 3.1 holds for $S = S_1 \cup \dots \cup S_n$.*
- (T) *Assume each S_i is compact in $\mathcal{T}_{\text{Zar}^*}$.*
- (I) *Assume that the S_i are pairwise incomparable.*

Then there exists $z \in K^\times$ with

$$v(z) = v(z_i) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n,$$

if and only if for any valuation w on K with a refinement in S_i and a refinement in S_j we have $w(z_i) = w(z_j)$.

Proof. It is clear that if such a z exists then every w with a refinement in S_i and a refinement in S_j satisfies $w(z_i) = w(z) = w(z_j)$. Conversely, if that compatibility condition is satisfied then the claim follows from Theorem 4.6 in Situation \mathfrak{m} with $x_i = z_i$ for all i . \square

Remark 5.2. It is worth pointing out in this context that any form of value approximation theorem like Corollary 5.1 implies the existence of a function ϕ as in Proposition 3.9, but of course it need in general not be given by a polynomial. It also implies the existence of elements b as in Lemma 4.2.

Remark 5.3. The assumption 5.1(T) cannot be replaced by compactness in the Zariski topology, as one can show with the following example in number fields – here an obstruction to value approximation is given by the class group.

Consider the number fields $K = \mathbb{Q}(\sqrt{-5})$; it is well-known that K has class number $h_K = 2$, with $\mathfrak{p}_0 = (2, 1 + \sqrt{-5})$ an example of a non-principal ideal, see for instance [Mar77, pp. 132–133]. The extension $L = K(\sqrt{-1})$ of K is of degree $2 = h_K$, and one can verify that it is unramified at all places. (Note that since $\mathbb{Q}(\sqrt{-1})/\mathbb{Q}$ is unramified over all finite primes except 2, it suffices to check that the primes of K above 2 are unramified in L .) Hence L is the Hilbert class field of K (see [Neu99, VI, Proposition 6.9]).

Consider the polynomial $f = X^2 - X - 1$, and let S be the set of valuations on K corresponding to prime ideals inert in L/K ; no such prime ideal has residue characteristic 5, since the prime ideal $(\sqrt{-5})$ is split in L/K . One verifies that L/K is generated by a zero of f , so in particular f is irreducible over K . The discriminant of f is 5. By standard results on the splitting of prime ideals in extensions, see [Neu99, I, Proposition 8.3], for any prime \mathfrak{p} of K inert in L (which necessarily does not contain 5), the reduction of f has no zero in $\mathcal{O}_K/\mathfrak{p}$. In particular, Assumption 3.1 is satisfied for the set S with the polynomial f and $t = 1$ by Example 3.5(1).

However, Corollary 5.1 does not transfer to this situation: Writing $S_1 = \{\mathfrak{p}_0\}$, $S_2 = S \setminus S_1$, we claim that there is no element $x \in K^\times$ with $v(x) = v(1 + \sqrt{-5})$ for $v \in S_1$ and $v(x) = v(1)$ for $v \in S_2$. If x were such an element, the ideal $(x)\mathfrak{p}_0^{-1}$ would be a product of prime ideals not inert in L/K . Since L/K is unramified of degree 2, it would be a product of prime ideals split in L/K , all of which are principal ideals in K by the theory of the Hilbert class field ([Neu99, VI, Corollary 7.4]). Hence \mathfrak{p}_0 itself would be principal, which is a contradiction. Therefore such an x cannot exist.

One can check that any Zariski open subset of S is empty or cofinite, hence the sets S_1 and S_2 are Zariski compact.

5.2. Residue approximation. The following ‘residue approximation’ appears for finitely many independent valuations already in [Kru31, Satz 17] (see also [Rib57, Lemme 6]) and for finitely many incomparable valuations in [Rib68, p. 143 Proposition 1], see also [Bou89, VI.7.2 Corollary 1] and [Efr06, Theorem 10.2.1]. A version for finitely many arbitrary valuations can be found in [Rib57, Lemme 11].

Corollary 5.4. *Let $S_1, \dots, S_n \subseteq S_{\text{val}}(K)$ and $x_1, \dots, x_n \in K$ with $x_i \in \mathcal{O}_v$ for each $v \in S_i$ and each i .*

- (U) *Assume that Assumption 3.1 holds for $S = S_1 \cup \dots \cup S_n$.*
- (T) *Assume each S_i is compact in $\mathcal{T}_{\text{Zar}^*}$.*
- (I) *Assume that the S_i are pairwise incomparable.*

Then there exists $x \in K$ with $x \in \mathcal{O}_v$ for each $v \in S_i$ for every i such that

$$\bar{x} = \bar{x}_i \text{ in } Kv \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Proof. This follows immediately from Theorem 4.6 in Situation \mathfrak{m} by choosing $z_i = 1$ for all i . \square

5.3. p -valuations. We now discuss approximation on the sets $S_\pi^e(K)$, which in particular includes the special case of p -valuations of bounded p -ramification index, cf. Example 3.6. We equip $S_\pi^e(K)$ with the constructible topology. Recall that $S_\pi^e(K)$ is compact and the topology coincides with the topology induced by $\mathcal{T}_{\text{Zar}^*}$ (Example 2.4). Let

$$R_\pi^e(K) = \bigcap_{v \in S_\pi^e(K)} \mathcal{O}_v$$

denote the corresponding holomorphy ring.

Corollary 5.5. *Let $S_1, \dots, S_n \subseteq S_\pi^e(K)$ be disjoint and closed, let $x_1, \dots, x_n \in K$, and let $z_1, \dots, z_n \in K^\times$. Assume that for any valuation w on K with a refinement in S_i and a refinement in S_j we have $w(x_i - x_j) \geq w(z_i) = w(z_j)$. Then there exists $x \in K$ with*

$$v(x - x_i) > v(z_i) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Proof. This follows from Theorem 4.6 in Situation **m** using Example 3.5(2). Note that two distinct valuations in $S_\pi^e(K)$ are always incomparable (Example 4.1). \square

Corollary 5.6. *Let $S_1, \dots, S_n \subseteq S_\pi^e(K)$ be disjoint and closed, $x_1, \dots, x_n \in R_\pi^e(K)$ and $k_1, \dots, k_n \in \mathbb{N}$. Then there exists $x \in K$ with*

$$v(x - x_i) > v(\pi^{k_i}) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Proof. This follows from Corollary 5.5: If w is a valuation with a refinement v_i in S_i and a refinement v_j in S_j , then $x_i, x_j \in R_\pi^e(K) \subseteq \mathcal{O}_{v_i} \subseteq \mathcal{O}_w$ and w is a proper coarsening of both v_i and v_j , hence $w(x_i - x_j) \geq 0 = w(\pi^{k_i}) = w(\pi^{k_j})$. \square

The argument for the following consequence is also contained in [Dar98, Propriété II.3.2].

Proposition 5.7. *For every $\pi \in K^\times$ and $e > 0$, the following statements are equivalent:*

- (1) *For every $v \in S_\pi^e(K)$, the holomorphy ring $R_\pi^e(K)$ is dense in \mathcal{O}_v in the v -topology.*
- (2) *The elements of $S_\pi^e(K)$ are pairwise independent.*

Proof. (1) \Rightarrow (2): Suppose two distinct $v_1, v_2 \in S_\pi^e(K)$ have a nontrivial common coarsening w . Let $a \in K^\times$ with $w(a) > 0$. As v_1 and v_2 are incomparable, Proposition 2.9 gives a $y \in K$ with $v_1(y) \geq v_1(\pi) > 0$ and $v_2(y) \leq v_2(\pi^{-1}) < 0$. If $R_\pi^e(K)$ was dense in \mathcal{O}_{v_1} , then there would exist $x \in K$ with $v_1(x) \geq 0$, $v_2(x) \geq 0$ and $v_1(x - y) > v_1(a)$. The latter condition implies that $w(x - y) \geq w(a) > 0$, but $v_2(y) < 0 \leq v_2(x)$ gives $v_2(x - y) < 0$ and thus $w(x - y) \leq 0$, a contradiction.

(2) \Rightarrow (1): Let $v_0 \in S_\pi^e(K)$, $y \in \mathcal{O}_{v_0}$ and $z \in K^\times$. We want to show there exists $x \in R_\pi^e(K)$ with $v_0(x - y) > v_0(z)$. Without loss of generality, $v_0(z) \geq 0$. Let

$$S_1 = \{v \in S_\pi^e(K) : v(y) \geq 0 \wedge v(z) \geq 0\}$$

and $S_2 = S_\pi^e(K) \setminus S_1$. Since in particular S_1 and S_2 are independent, we can apply Corollary 5.5 to $x_1 = y$, $x_2 = 0$, $z_1 = z$, $z_2 = 1$ to get $x \in K$ with $v(x - y) > v(z) \geq 0$ for $v \in S_1$, and $v(x) \geq 0$ for $v \in S_2$. In particular $v_0(x - y) > v_0(z)$ and $x \in R_\pi^e(K)$. \square

Example 5.8. An example where Proposition 5.7 can be applied is when K is a so-called pseudo p -adically closed field: If we set $\pi = p$ and $e = 1$, then $S_\pi^e(K)$ equals the set of p -valuations of p -rank 1, and any two of these are independent, see Theorem C and the remark following Proposition D of [GJ91].

5.4. Comparison with strong approximation. We now want to compare our approximation theorems with the well known results for a global field K . Beyond the Weak Approximation Theorem 1.1 valid for any field, in global fields we have the following stronger result.

Theorem 5.9 (Strong Approximation, [CF67, Chapter II §15 Theorem]). *Let K be a global field and $S \subsetneq S_{\text{abs}}(K) \setminus \{v_{\text{trivial}}\}$. For each $v \in S$, let $x_v \in K$ and $\epsilon_v > 0$ be given such that $x_v \in \mathcal{O}_v$ and $\epsilon_v = 1$ for almost all $v \in S \cap S_{\text{val}}(K)$. Then there exists an $x \in K$ with $|x - x_v|_v \leq \epsilon_v$ for all $v \in S$.*

As usual in algebraic number theory, in this situation a nontrivial element of $S_{\text{abs}}(K)$ is called a place. The condition that S excludes at least one place can clearly not be omitted due to the product formula (e.g. see [Neu99, III, (1.3)]) – for instance, there is no element of K^\times which is of norm ≤ 1 at all places and of norm < 1 at one of them.

To compare with our theorems, one first has to analyse the topologies on S defined in Section 2. One easily checks that the Zariski topology on S is exactly the cofinite topology; in particular any subset of S is Zariski compact. On the other hand, the Hochster dual of the Zariski topology is the discrete topology, so S is never compact unless it is finite. (It is important here that we excluded the trivial valuation from S .)

This means that while our approximation theorem in Situation \mathfrak{m} is quite weak for global fields – we can only approximate on finite sets S_i , i.e. do not obtain anything stronger than the weak approximation Theorem 1.1 –, we can use our approximation theorem in Situation \mathcal{O} to recover strong approximation under additional hypotheses.

Proof of Theorem 5.9 when S contains no complex places and Assumption 3.1 holds. For each $v \in S$ let x_v and ϵ_v be given. Write $S_0 \subseteq S$ for the set of finite places $v \in S$ with $x_v = 1$, $\epsilon_v = 1$. The set $S \setminus S_0$ is finite, so enumerate it as $\{v_1, \dots, v_n\}$, and for each i find a $z_i \in K^\times$ with $|z_i|_{v_i} \leq \epsilon_i$. Writing $z_0 = x_0 = 1$ and $S_i = \{v_i\}$, $x_i = x_{v_i}$ for $i \geq 1$, we apply Theorem 4.6 in Situation \mathcal{O} to the sets S_i and elements x_i, z_i for $i = 0, \dots, n$. The element x thus obtained is as desired. \square

By Example 3.8, this proves strong approximation in some situations of sets of places S with density arbitrarily close to 1, i.e. we only have to exclude a set of places of small density. We will see how to lift the prohibition on complex places in Section 6. However, we cannot reach the full statement of strong approximation, in which only a single place needs to be omitted, since for any non-constant $f \in K[X]$ and $t \in K^\times$ the Chebotarev Density Theorem shows that there is always a positive density of finite places v of K with $v(t) = 0$ in whose residue field the reduction of f has a zero, hence violating Assumption 3.1.

5.5. Kronecker dimension one and reduction to finitely generated fields. For so-called fields of Kronecker dimension one, i.e. algebraic extensions of \mathbb{Q} and algebraic extensions of some rational function field $\mathbb{F}_q(T)$, most of our approximation results, or some variants of it, are very easy to prove since they can be reduced to approximation results in finite extensions of \mathbb{Q} respectively $\mathbb{F}_q(T)$. For example, one even has the following stronger result:

Proposition 5.10. *Let K be a field of Kronecker dimension one, $S_0, \dots, S_n \subseteq S(K)$ pairwise disjoint nonempty constructibly closed sets, $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$. Then there exists $x \in K$ with*

$$x - x_i \in z_i \mathcal{O}_v \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Proof. As $S(K)$ is the inverse limit of $S(K_0)$ for the finitely generated subfields K_0 of K , and $S_{\text{abs}}(K_0) \setminus S_{\text{val}}(K_0)$ is finite for all of these, the space $S(K)$ is compact Hausdorff and therefore normal. We can thus assume without loss of generality that the S_i are open-closed (and still pairwise disjoint), i.e. of the form

$$S_i = \{v \in S(K) : a_{i1}, \dots, a_{ik_i} \in \mathcal{O}_v, b_{i1}, \dots, b_{il_i} \in \mathfrak{m}_v\}$$

with elements $a_{ij}, b_{ij} \in K$. The subfield K_0 of K generated by all a_{ij}, b_{ij}, x_i, z_i is then without loss of generality a global field, and

$$S'_i := \{v \in S(K_0) : a_{i1}, \dots, a_{ik_i} \in \mathcal{O}_v, b_{i1}, \dots, b_{il_i} \in \mathfrak{m}_v\}$$

consists exactly of the restrictions of the elements of S_i to K_0 . In particular, S'_0, \dots, S'_n are again nonempty and pairwise disjoint. Now let

$$T'_i = \{v \in S'_i : x_i \in \mathcal{O}_v, z_i^{-1} \in \mathcal{O}_v\}.$$

Then $S'_i \setminus T'_i$ is finite for every i , so by Theorem 5.9 there exists $x \in K_0$ such that $x - x_i \in z_i \mathcal{O}_v$ for every $v \in S'_i \setminus T'_i$ and $x \in \mathcal{O}_v$ for every $v \in T'_i$, for $i = 1, \dots, n$. This x then satisfies $x - x_i \in z_i \mathcal{O}_v$ for every $v \in S_i$ and every $i = 1, \dots, n$. \square

Note that this proposition needs neither a (U) condition, since the set of localities on the global field K_0 has the property that every $a \in K_0$ lies in \mathcal{O}_v^\times for almost all $v \in S(K_0)$, nor an (I) condition, since localities on a global field are automatically pairwise independent.

This might suggest that one can possibly reduce Theorem 4.6 to the special case where the field is finitely generated over its prime field, by replacing the general field K by a suitable finitely generated subfield K_0 . This however does not seem to be of much use, mainly since in general neither of the two properties of global fields named in the previous paragraph holds for K_0 ; in fact, even if a set $S \subseteq S(K)$ satisfies some independence or compatibility condition, the restriction of S to K_0 need not.

5.6. The (U) condition. We have seen in Remarks 4.9 and 5.3 that in our theorems the conditions (I) and (T) cannot be dropped or significantly weakened. It remains to justify condition (U), i.e. essentially Assumption 3.1, which is technical and may appear unnatural.

We have seen above that the product formula alone may necessitate the omission of some place. The following example justifies our stronger assumption. We focus on a situation with only valuations, so that only conditions 3.1(i, ii, iii) play a role.

Example 5.11. Let P be a set of prime numbers such that for every number field L there exists a $p \in P$ (or equivalently infinitely many $p \in P$) such that the prime ideal (p) is completely split in L/\mathbb{Q} . We may of course simply take P to be the set of all prime numbers.

We now consider $K = \mathbb{Q}(T)$; we will construct two Zariski compact sets S_1, S_2 of valuations on K , both only consisting of p -valuations (of p -rank 1) for some $p \in P$, such that a certain approximation problem has no solution. We let S_2 consist of a single valuation, namely the refinement of the degree valuation on K by the q -adic valuation on \mathbb{Q} for some fixed $q \in P$. To construct S_1 , first fix an enumeration f_1, f_2, \dots of the irreducible monic polynomials in $\mathbb{Q}[T]$ and an enumeration x_1, x_2, \dots of the non-zero elements of K . To each f_i there is an associated discrete valuation v_{f_i} on K , trivial on \mathbb{Q} .

For each i , construct a valuation on K in the following way. The residue field of v_{f_i} is a finite extension of \mathbb{Q} , so by assumption on P it carries a p -valuation v_p of p -rank 1 for some $p \in P$. We may even choose p such that for all x_j , $1 \leq j \leq i$, with $v_{f_i}(x_j) = 0$ we have $(v_p \circ v_{f_i})(x_j) = 0$, as this latter condition only excludes finitely many p . We let $v_i = v_p \circ v_{f_i}$, and take $S_1 = \{v_i : i \geq 1\}$.

We claim that S_1 , as a subspace of $S_{\text{val}}(K)$ with the Zariski topology, carries the cofinite topology. To see this, observe that for every $x_j \in K^\times$, we have $v_i(x_j) = 0$ unless either $i < j$ or $v_{f_i}(x_j) \neq 0$, each of which only happens for finitely many i . Hence every nonempty Zariski-open set is cofinite, and in particular S_1 is compact. Since the valuations in S_1 are pairwise incomparable, the topology is T_1 (Remark 2.6) and therefore the Zariski topology is exactly the cofinite topology.

Consider now the following approximation problem: We demand an $x \in K$ such that $v(x - 0) \geq v(1)$ for all $v \in S_1$, and $w(x - T^{-1}) \geq w(T^{-2})$ for $w \in S_2$. Such an x would be integral at all v_{f_i} and furthermore integral at the degree valuation, hence necessarily constant. However, then $w(x - T^{-1}) \geq w(T^{-2})$ would be violated. Therefore this approximation problem is not solvable, in spite of S_1 and S_2 being compact sets of valuations, any two of which are independent.

Note that if P did not satisfy our initial condition, i.e. if there exists a number field L/\mathbb{Q} such that for no $p \in P$ is the ideal (p) completely split in L/\mathbb{Q} , then we may as well enlarge L to a totally imaginary Galois extension of \mathbb{Q} ; in this situation, Example 3.8 (where we choose g to be the minimal polynomial of an integral primitive element of L/\mathbb{Q}) shows that Assumption 3.1 applies with $t \in \mathbb{Q}^\times$ (or even $t = 1$ since we are not interested in orderings), and therefore Theorem 4.6 is applicable to sets S consisting only of valuations with residue field \mathbb{F}_p for

some $p \in P$ unramified in L/\mathbb{Q} . Hence Example 5.11 shows that condition 4.6(U) cannot be substantially weakened in Situation \mathcal{O} .

5.7. Affine families of valuations. We now briefly discuss the relation between our results and the approximation results in the work of Ershov, e.g. [Ers94, Ers01, Ers06]. One of the most general results Ershov obtains is the following, which we have paraphrased.

Theorem 5.12 (see [Ers01, Proposition 2.6.2.]). *Let $S_1, \dots, S_n \subseteq S_{\text{val}}(K) \setminus \{v_{\text{trivial}}\}$ pairwise disjoint, and let $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$. Write $S = S_1 \cup \dots \cup S_n$.*

(U) *Assume that $R = \bigcap_{v \in S} \mathcal{O}_v$ is a Prüfer ring with quotient field K .¹²*

(T) *Assume that each S_i is compact in the Zariski topology.*

(I) *Assume that the elements of S are pairwise independent.*

Then there exists $x \in K$ with

$$v(x - x_i) \geq v(z_i) \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Condition 5.12(U) is satisfied for example in the situation of Example 3.5(2), see [Ers94, Proposition 3]. In particular, Theorem 1.5 follows from Theorem 5.12 applied to $S_\pi^1(K)$. Condition 5.12(U) is also satisfied in the situation of Example 3.5(1), see [Ers01, Proposition 2.3.3] or [Roq73, Theorem 1].

On the other hand, Theorem 5.12 can be proven by our methods under the stronger assumption that R is not only Prüfer but satisfies the condition explained in Remark 3.12.

6. ADDING FINITELY MANY EXCEPTIONAL LOCALITIES

The standing Assumption 3.1 requires a polynomial $f \in K[X]$ and an element $t \in K^\times$ such that $f(x) \notin t\mathfrak{m}_v$ for all $x \in K$ and all localities v under consideration. If v is a rank-1 valuation, this means in particular that f has no zero in the completion of K with respect to v , so we cannot hope to cover rank-1 valuations with algebraically closed completion. For the same reason, our method as is cannot cover absolute values with completion \mathbb{C} , although this is desirable for analogy with Theorem 1.1. It turns out, however, that at least finitely many such exceptional localities can be added to our main theorem.

The following lemma is a variant of Proposition 2.9.

Lemma 6.1. *Let $v_0, v_1, \dots, v_n \in S(K)$ and $z_0, \dots, z_n \in K^\times$ such that for any valuation w coarsening v_0 and some v_i , $i > 0$, we have $w(z_0) \geq w(z_i)$.*

(1) *If every v_i with $i > 0$ is strongly incomparable to v_0 , then there exists $z \in K^\times$ with $z \in z_i\mathfrak{m}_{v_i}$ for all $i > 0$ and $z^{-1} \in z_0^{-1}\mathfrak{m}_{v_0}$.*

(2) *If every v_i with $i > 0$ is either strongly incomparable to or a proper refinement of v_0 , then there exists $z \in K^\times$ with $z \in z_i\mathfrak{m}_{v_i}$ for all $i > 0$ and $z^{-1} \in z_0^{-1}\mathcal{O}_{v_0}$.*

Proof. Note that we may freely scale the z_i by a common factor, so assume without loss of generality that $z_0 = 1$.

Consider (1). Assume first that v_0, \dots, v_n are dependent, say with nontrivial finest common coarsening w . Any z_i with $w(z_i) < 0$ may be replaced by 1, since this only strengthens the conclusion. Hence we have $w(z_i) = 0$ for all i , and we may reduce to a problem in the residue field Kw , on which v_0, \dots, v_n induce localities $\bar{v}_0, \dots, \bar{v}_n$: Any lift $z \in K^\times$ of $\bar{z} \in (Kw)^\times$ satisfying $\bar{z} \in \bar{z}_i\mathfrak{m}_{\bar{v}_i}$ for all $i > 0$ and $\bar{z}^{-1} \in \bar{z}_0^{-1}\mathfrak{m}_{\bar{v}_0}$ is as desired. Hence we have reduced to a problem in the residue field, where the $\bar{v}_0, \dots, \bar{v}_n$ are not all dependent. Therefore let us assume henceforth that v_0, \dots, v_n are not all dependent.

Inductively, we may first solve the problem restricted to all v_i which are dependent with v_0 , so let $z' \in K^\times$ with $z'^{-1} \in z_0^{-1}\mathfrak{m}_{v_0}$ and $z' \in z_i\mathfrak{m}_{v_i}$ for all v_i dependent with v_0 . Note that these

¹²In [Ers01, Proposition 2.6.2.], the condition is that S is *affine*, which he shows to be the case iff S satisfies (U), is compact in the Zariski topology and consists of pairwise incomparable valuations, see [Ers01, Proposition 2.3.4, Corollary 2.3.2]. The latter two are implied by (T) and (I).

conditions are then also satisfied in a v_0 -neighbourhood of z' . Likewise, for any v_i independent from v_0 there is a v_i -open set of z'_i such that $z'_i \in z_j \mathfrak{m}_{v_j}$ for all v_j dependent with v_i . Then Theorem 2.8 gives z as desired.

For (2), if v_0 is strongly incomparable to every other v_i , then we may solve the stronger problem (1), so assume this is not the case, i.e. some v_i properly refines v_0 . In particular v_0 is a valuation. Since the residue field of v_0 carries a nontrivial locality, it carries infinitely many pairwise independent valuations, so we may pick a refinement v'_0 of v_0 with $v'_0 \vee v_i = v_0$ for any of the v_i refining v_0 . Note that any w coarsening v'_0 and some v_i will also coarsen $v_0 = v'_0 \vee v_i$, and hence satisfies $w(z_0) \geq w(z_i)$. We can then solve problem (1) for v'_0, v_1, \dots, v_n , and any solution thereof is as desired. \square

The following results are again to be understood in the two situations \mathcal{O} and \mathfrak{m} as before. This first proposition is an extension of Theorem 1.3 to include orderings and complex absolute values.

Proposition 6.2. *Let $n > 1$, $v_1, \dots, v_n \in \mathcal{S}(K)$, $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$. Assume that whenever w coarsens v_i and v_j , $i \neq j$, then $x_i - x_j \in z_i \mathcal{O}_w = z_j \mathcal{O}_w$. Furthermore assume in Situation \mathfrak{m} that for any i, j with $x_i \neq x_j$ the localities v_i and v_j are strongly incomparable; in Situation \mathcal{O} , assume the same only when v_i and v_j are orderings. Then there exists $x \in K$ with $x - x_i \in z_i \mathfrak{B}_{v_i}$ for all i .*

Proof. In Situation \mathcal{O} , if v_i refines v_j , then $x_i + z_i \mathfrak{B}_{v_i} \subseteq x_j + z_j \mathfrak{B}_{v_j}$, so we may simply remove v_j from the list. We can repeat this until the v_i are pairwise incomparable. If only one v_i remains, the problem has become trivial, otherwise all conditions for the stronger result in Situation \mathfrak{m} are satisfied, so it suffices to consider this situation.

We use induction on n (where K varies across the class of all fields). If the v_i are pairwise independent the statement follows from Theorem 2.8. If the v_i are all dependent with nontrivial finest common coarsening w , we may assume after scaling and shifting as in Remark 2.10 that $w(z_i) = 0$ and $x_i \in \mathcal{O}_w$ for all i . It then suffices to solve the induced problem in the residue field Kw , i.e. to find $\bar{x} \in Kw$ with $\bar{x} - \bar{x}_i \in \bar{z}_i \mathfrak{m}_{\bar{v}_i}$ for all i , since any lift $x \in K$ of \bar{x} will be as desired. Solving the induced problem in Kw is possible inductively, since for any v_i and v_j which induce the same locality on the residue field (i.e. are in particular not strongly incomparable), we have assumed that $x_i = x_j$, so we obtain only one condition in the residue field for v_i and v_j .

Therefore assume that the v_i are neither pairwise independent nor all dependent. After reordering, let v_1, \dots, v_k with $1 < k < n$ be a maximal dependent subset with nontrivial common coarsening w . By scaling and shifting we may assume that $w(z_i) = 0$ and $x_i \in \mathcal{O}_w$ for all $i \leq k$. We can apply the induction hypothesis in the residue field of w to obtain an $\bar{x} \in Kw$ which satisfies $\bar{x} - \bar{x}_i \in \bar{z}_i \mathfrak{m}_{\bar{v}_i}$ for any $i \leq k$. Take any lift $x \in K$ of \bar{x} . For $i \leq k$ we then have $x + z_i \mathfrak{m}_w \subseteq x_i + z_i \mathfrak{m}_{v_i}$. Hence we may replace the conditions with respect to the v_1, \dots, v_k by a single condition with respect to w , reducing the number of conditions. Using the induction hypothesis once more proves the claim. \square

Theorem 6.3. *Let $S_1, \dots, S_n \subseteq \mathcal{S}(K)$, $x_1, \dots, x_n \in K$ and $z_1, \dots, z_n \in K^\times$.*

- (U) *Assume that there exist $f \in K[X]$ and $t \in K^\times$ such that conditions 3.1(i, iii, iv) are satisfied for all localities in all S_i , and condition 3.1(ii) is satisfied for all localities in all infinite S_i .*
- (T) *Assume that each S_i is compact in the given topology.*
- (I) *Assume that for any valuation w on K with a refinement in S_i and a refinement in S_j we have $w(x_i - x_j) \geq w(z_i) = w(z_j)$; assume further that the S_i are pairwise t -incomparable, and in Situation \mathfrak{m} furthermore pairwise incomparable.*

Then there exists $x \in K$ with

$$x - x_i \in z_i \mathfrak{B}_v \text{ for all } v \in S_i, \text{ for } i = 1, \dots, n.$$

Proof. We may assume that all S_i are nonempty. After reordering if necessary, say S_1, \dots, S_k are infinite and S_{k+1}, \dots, S_n are finite. In Situation \mathcal{O} , we may furthermore assume that no

element of S_i with $i > k$ is a coarsening of an element of S_j with $j \leq k$, since such a coarsening could simply be removed.

We first apply Proposition 6.2 to the elements of S_{k+1}, \dots, S_n to obtain an element $x' \in K$ such that $x' - x_i \in tz_i \mathfrak{B}_v$ for all $v \in S_i$, $i > k$. We secondly apply Theorem 4.6 to the sets S_1, \dots, S_k , elements x_1, \dots, x_k and tz_1, \dots, tz_k to obtain an element $x'' \in K$ with $x'' - x_i \in tz_i \mathfrak{B}_v$ for all $v \in S_i$, $i \leq k$.

We want to find $x \in K$ with $x - x' \in tz_i \mathfrak{B}_v$ for $v \in S_i$, $i > k$, and $x - x'' \in tz_i \mathfrak{B}_v$ for $v \in S_i$, $i \leq k$; such an x is as desired, by Assumption 3.1(iv). Note that we have the basic compatibility condition

$$w(x' - x'') \geq \min\{w(x_i - x'), w(x_i - x_j), w(x_j - x'')\} \geq w(tz_i) = w(tz_j)$$

for any valuation w coarsening elements of S_i and S_j , $i > k \geq j$.

If $x' = x''$, we set $x = x'$, and in the remaining case we may assume without loss of generality that $x' = 1$ and $x'' = 0$, by scaling and shifting all x_i and x' and x'' , and scaling all z_i . It now suffices to find $b \in K^\times$ with $b \in t\mathfrak{m}_v \cap tz_i \mathfrak{m}_v$ for $v \in S_i$, $i > k$, and $b^{-1} \in t\mathfrak{B}_v \cap tz_i \mathfrak{B}_v$ for all $v \in S_i$, $i \leq k$; with such b we can apply Lemma 4.3 to obtain x . (Note that by Remark 4.4 only condition (iii) is necessary for $v \in S_i$, $i > k$, to apply the lemma.)

To find such b , we imitate the proof of Lemma 4.2. For any $i \leq k$ and $v' \in S_i$, there exists an element $b_{i,v'} \in K^\times$ with $b_{i,v'} \in t\mathfrak{m}_v \cap tz_j \mathfrak{m}_v$ for all $v \in S_j$, $j > k$, and $b_{i,v'}^{-1} \in t\mathfrak{B}_{v'} \cap tz_i \mathfrak{B}_{v'}$; this follows from Lemma 6.1 applied to v' and the $v \in S_{k+1} \cup \dots \cup S_n$ with an element $z_{v'} \in K^\times$ chosen to satisfy $z_{v'} \mathfrak{B}_{v'} = t\mathfrak{B}_{v'} \cap tz_i \mathfrak{B}_{v'}$ and elements $z_v \in K^\times$ chosen to satisfy $z_v \mathfrak{m}_v = t\mathfrak{m}_v \cap tz_j \mathfrak{m}_v$ for any $j > k$ with $v \in S_j$ (note that if $v \in S_j \cap S_{j'}$ then $z_j \mathfrak{m}_v = z_{j'} \mathfrak{m}_v$ by (I)). Note that if w coarsens v' and some $v \in S_j$, then $w(z_i) = w(z_j)$, hence $w(z_{v'}) = w(z_v)$.

By compactness of the S_i , there exists a finite list P of pairs (i, v) such that for any $i \leq k$ and $v' \in S_i$ there exists $v \in S_i$ with $(i, v) \in P$ and $b_{i,v}^{-1} \in t\mathfrak{B}_{v'} \cap tz_i \mathfrak{B}_{v'}$. Now take $b = \phi(b_1, \dots, b_m)$, where b_1, \dots, b_m are the $\{b_{i,v} : (i, v) \in P\}$ in arbitrary order. By Corollary 3.10, the first part of which only requires conditions (i, iii), this b is as desired. \square

Note that imposing conditions (i, iii, iv) on those S_i which are finite is quite weak in practice. For any complex absolute value $|\cdot|$, the conditions are satisfied for example for $f = \sum_{i=0}^d a_i X^i \in K[X]$ and $t \in K^\times$ if

$$\sum_{i=0}^d |a_i| \leq 1, \quad |t| + \sum_{i=0}^{d-1} |a_i| \leq |a_d|, \quad \text{and} \quad |t| \leq \frac{1}{2},$$

which are conditions we have already seen for orderings in Example 3.7. In particular, Example 3.8 extends to also cover complex absolute values in this way.

For a valuation v , conditions (i, iii, iv) are satisfied for example if f has coefficients in \mathcal{O}_v and $v(t) \geq v(a_d) = 0$.

7. APPROXIMATION OF VALUES OF RATIONAL FUNCTIONS

Let $S \subseteq S(K)$ be nonempty. We write

$$\mathfrak{m}_S := \bigcap_{v \in S} \mathfrak{m}_v.$$

Note that $\mathfrak{m}_S \cdot \mathfrak{m}_S \subseteq \mathfrak{m}_S$, although \mathfrak{m}_S is never an element of $\mathcal{B}(K)$ as defined in Section 2. Let us assume throughout that S does not contain the trivial valuation, satisfies Assumption 3.1, and is compact in $\mathcal{T}_{\text{Zar}^*}$. Note that $S \subseteq S_{\text{val}}(K) \cup S_{\text{ord}}(K)$ since complex places always violate Assumption 3.1.

Lemma 7.1. *Then $\mathfrak{m}_S \neq \{0\}$, and the family*

$$\mathcal{M}_S := \{z\mathfrak{m}_S \mid z \in \mathfrak{m}_S \setminus \{0\}\}$$

is a filter base.

Proof. First, a simple compactness argument shows that $\mathfrak{m}_S \neq \{0\}$, as follows. Each $v \in S$ is nontrivial, so there exists $z_v \in \mathfrak{m}_v \setminus \{0\}$. By compactness in $\mathcal{T}_{\text{Zar}^*}$, there are finitely many $v_1, \dots, v_n \in S$ such that

$$S \subseteq \bigcup_{i=1}^n \{v \in S(K) : z_{v_i} \in \mathfrak{m}_v\}.$$

Therefore $\bigcap_{i=1}^n z_{v_i} \mathcal{O}_v \subseteq \mathfrak{m}_v$, for all $v \in S$. Choosing ϕ as in Corollary 3.10, we let $z_0 = \phi(z_{v_1}^{-1}, \dots, z_{v_n}^{-1})^{-1}$, which is nonzero. By Corollary 3.10(2), for each $v \in S$, $z_0 \in \bigcap_{i=1}^n z_{v_i} \mathcal{O}_v$. Therefore $z_0 \in \mathfrak{m}_S \setminus \{0\}$, and so \mathcal{M}_S is a nonempty family of nonempty sets.

To see that \mathcal{M}_S is a filter base, we let $z_1, z_2 \in \mathfrak{m}_S \setminus \{0\}$. Let $y := \phi(z_1^{-1}, z_2^{-1})^{-1}$, which we note is nonzero. By Proposition 3.9(2) we have $y \in z_1 \mathcal{O}_v \cap z_2 \mathcal{O}_v \subseteq \mathfrak{m}_v$ for each $v \in S$. Thus $y \mathfrak{m}_S \in \mathcal{M}_S$ and $y \mathfrak{m}_S \subseteq y \mathfrak{m}_v \subseteq z_1 \mathfrak{m}_v \cap z_2 \mathfrak{m}_v$ for every $v \in S$, and therefore $y \mathfrak{m}_S \subseteq z_1 \mathfrak{m}_S \cap z_2 \mathfrak{m}_S$, as required. \square

Consequently, there is a filter $\mathcal{N}_{S,0}$ of which \mathcal{M}_S is a filter base.

Lemma 7.2. *Let \mathcal{F} be a filter on K . Then there exists a Hausdorff field topology \mathcal{T} on K such that \mathcal{F} is the filter of \mathcal{T} -neighbourhoods of 0 if and only if the following conditions hold.*

- (1) $\forall x \in K^\times \exists V \in \mathcal{F} : x \notin V$
- (2) $\forall U \in \mathcal{F} \exists V \in \mathcal{F} : V + V \subseteq U$
- (3) $\forall U \in \mathcal{F} \exists V \in \mathcal{F} : V \subseteq -U$
- (4) $\forall U \in \mathcal{F} \exists V \in \mathcal{F} : V \cdot V \subseteq U$
- (5) $\forall U \in \mathcal{F} \forall x \in K^\times \exists V \in \mathcal{F} : xV \subseteq U$
- (6) $\forall U \in \mathcal{F} \exists V \in \mathcal{F} : (1 + V)^{-1} \subseteq 1 + U$

Proof. By [War89, Ch. II Thm. 11.4], \mathcal{F} satisfies (2)-(5) if and only if \mathcal{F} is the filter of \mathcal{T} -neighborhoods of 0 for a ring topology \mathcal{T} , which is then uniquely determined by \mathcal{F} . In this case, by continuity of addition, the sets $1 + U$ form the filter of \mathcal{T} -neighborhoods of 1, and therefore (6) is equivalent to the continuity of inversion at 1. By continuity of multiplication, this is already equivalent to the continuity of inversion on K^\times . Finally, for the filter of \mathcal{T} -neighborhoods of 0, (1) is equivalent to $\bigcap \mathcal{F} = \{0\}$, which in any topological group holds if and only if the topology is Hausdorff ([War89, Ch. I Thm. 1.7]). \square

Lemma 7.3. *The filter $\mathcal{N}_{S,0}$ satisfies conditions (1)-(6) of Lemma 7.2.*

Proof. It suffices to verify (1)-(6) for the filter base \mathcal{M}_S . We choose $z_0 \in \mathfrak{m}_S \setminus \{0\}$ as in Lemma 7.1 above. Let $x \in K^\times$ and let $y_0 := \phi(x^{-1}, z_0^{-1})^{-1}$, which we note is nonzero. By Proposition 3.9(2), we have $y_0 \in x \mathcal{O}_v \cap z_0 \mathcal{O}_v \subseteq \mathfrak{m}_v$, for all $v \in S$. In particular $y_0 \in \mathfrak{m}_S \setminus \{0\}$, and $x \notin y_0 \mathfrak{m}_S$. This proves (1).

Turning to the other conditions, let $z_1 \in \mathfrak{m}_S \setminus \{0\}$. Then $y_1 := z_1 t \in \mathfrak{m}_S \setminus \{0\}$. It follows from Assumption 3.1(iv) that $\mathfrak{m}_S + \mathfrak{m}_S \subseteq t^{-1} \mathfrak{m}_S$. Then

$$y_1 \mathfrak{m}_S + y_1 \mathfrak{m}_S \subseteq z_1 \mathfrak{m}_S,$$

which verifies (2). Condition (3) holds because $-\mathfrak{m}_S = \mathfrak{m}_S$.

Let $z_2 \in \mathfrak{m}_S \setminus \{0\}$, and simply choose $y_2 = z_2$. Then

$$(y_2 \mathfrak{m}_S) \cdot (y_2 \mathfrak{m}_S) = z_2^2 \mathfrak{m}_S \cdot \mathfrak{m}_S \subseteq z_2 \mathfrak{m}_S,$$

which verifies (4).

Let $z_3 \in \mathfrak{m}_S \setminus \{0\}$ and let $x \in K^\times$. We let $y_3 := z_3 \cdot \phi(x, 1)^{-1}$, which we note is nonzero. By Proposition 3.9(2), we have $y_3 \in z_3 x^{-1} \mathcal{O}_v \cap z_3 \mathcal{O}_v$, for all $v \in S$. Therefore $y_3 \in \mathfrak{m}_S \setminus \{0\}$, since $z_3 \in \mathfrak{m}_S$. Finally we have $x y_3 \mathfrak{m}_S \subseteq z_3 \mathfrak{m}_S$, which verifies (5).

Finally, let $z_4 \in \mathfrak{m}_S$, $z_4 \neq 0$. For any valuation $v \in S$, we have $(1 + z_4 \mathfrak{m}_v)^{-1} \subseteq 1 + z_4 \mathfrak{m}_v$. For any ordering $v \in S$, we have $(1 + \frac{z_4}{2} \mathfrak{m}_v)^{-1} \subseteq 1 + z_4 \mathfrak{m}_v$ since for any real number $r \in (0, 1)$ we have $1 - r \leq (1 + \frac{r}{2})^{-1} < (1 - \frac{r}{2})^{-1} \leq 1 + r$. Let $y := \phi(z_4^{-1}, 2z_4^{-1})^{-1}$. As above, we have $y \mathfrak{m}_S \in \mathcal{M}_S$ and $y \mathfrak{m}_S \subseteq z_4 \mathfrak{m}_S \cap \frac{z_4}{2} \mathfrak{m}_S$. Then $(1 + y \mathfrak{m}_S)^{-1} \subseteq 1 + z_4 \mathfrak{m}_S$, which verifies (6). \square

It follows from Lemma 7.3 that there is a (unique) field topology on K of which $\mathcal{N}_{S,0}$ is the filter of neighbourhoods of 0. We call this topology the S -topology. A subset of K which is open in the S -topology is said to be S -open. An S -ball is a set of the form

$$B_S(x, z) := x + z\mathfrak{m}_S,$$

for $x \in K$ and $z \in K^\times$.

Lemma 7.4. *All S -balls are S -open.*

Proof. It suffices to show that \mathfrak{m}_S is S -open. Let $x \in \mathfrak{m}_S$. We must find $z \in \mathfrak{m}_S \setminus \{0\}$ such that $x + z\mathfrak{m}_S \subseteq \mathfrak{m}_S$. First choose any $z_0 \in \mathfrak{m}_S \setminus \{0\}$, and write $z := z_0\phi((1-x)^{-1}, (1+x)^{-1})^{-1}$, and note that z is non-zero. For each $v \in S$, either $1-x$ or $1+x$ is in \mathcal{O}_v , and therefore $z \in z_0(1-x)\mathcal{O}_v \cap z_0(1+x)\mathcal{O}_v \subseteq \mathfrak{m}_v$, by Proposition 3.9(2). In particular $z \in \mathfrak{m}_S \setminus \{0\}$.

If $v \in S$ is a valuation then certainly $x + z\mathfrak{m}_v \subseteq \mathfrak{m}_v$. On the other hand, suppose that $v \in S$ is an ordering and let $y \in \mathfrak{m}_v$. If $0 \leq_v x <_v 1$, then $1-x \leq_v 1+x$, so $x-1 <_v z <_v 1-x$. Otherwise if $-1 <_v x <_v 0$, then $-x-1 <_v z <_v 1+x$. In either case we have $x+zy \in \mathfrak{m}_v$, and so $x + z\mathfrak{m}_v \subseteq \mathfrak{m}_v$, for all orderings $v \in S$. Therefore $x + z\mathfrak{m}_S \subseteq \mathfrak{m}_S$, as required. \square

It follows from Lemma 7.4 that the family

$$\{B_S(x, z) : x \in K, z \in \mathfrak{m}_S \setminus \{0\}\}$$

is a base for the S -topology. Note that the S -topology is finer than the v -topology, for every $v \in S$. For example, if $v \in S$ is a valuation, then

$$\mathfrak{m}_v = \bigcup_{x \in \mathfrak{m}_S} x + \mathfrak{m}_S.$$

If $S = \{v\}$ is a singleton, then $B_v(x, z) = B_{\{v\}}(x, z)$, and thus the S -topology coincides with the v -topology. In this case we will speak of the S -topology and S -balls to mean the v -topology and v -balls, whether or not Assumption 3.1 is satisfied.

For each $m \in \mathbb{N}$, we define the S -topology on K^m to be the product topology induced by the S -topology on K . An S -ball in K^m is a set of the form

$$B_S(\underline{x}, \underline{z}) := \prod_{i=1}^m B_S(x_i, z_i),$$

for tuples $\underline{x} = (x_1, \dots, x_m) \in K^m$ and $\underline{z} = (z_1, \dots, z_m) \in (K^\times)^m$. For consistency of notation, we write $B_v(\underline{x}, \underline{z}) := B_{\{v\}}(\underline{x}, \underline{z})$. For $D \subseteq K^l$, a function $f : D \rightarrow K^m$ is said to be S -continuous if it is continuous with respect to the S -topologies. Furthermore, f is S -hereditarily continuous if it is S' -continuous for each nonempty $\mathcal{T}_{\text{Zar}^*}$ -compact $S' \subseteq S$.

Example 7.5. If $g \in K(x_1, \dots, x_l)^m$ is a tuple of rational functions given by $g_i = \frac{h_i}{k_i}$ with $h_i, k_i \in K[x_1, \dots, x_l]$ coprime, then the domain

$$D_g = \{\underline{x} \in K^l : k_i(\underline{x}) \neq 0 \text{ for all } i\}$$

of g is open with respect to each $v \in S$, and the induced map $g : D_g \rightarrow K^m$ is S -hereditarily continuous, since each of the S' -topologies (for $S' \subseteq S$ $\mathcal{T}_{\text{Zar}^*}$ -compact) is a field topology.

Proposition 7.6. *Let $S_1, \dots, S_n \subseteq S(K) \setminus \{v_{\text{trivial}}\}$ be nonempty.*

- (U) *Assume that there exist $f \in K[X]$ and $t \in K^\times$ such that conditions 3.1(i, iii, iv) are satisfied for all localities in all S_i , and condition 3.1(ii) is satisfied for all localities in all S_i with $|S_i| > 1$.*
- (T) *Assume each S_i is compact in $\mathcal{T}_{\text{Zar}^*}$.*
- (I) *Assume that the S_i are pairwise independent.*

For each i , let $A_i \subseteq K^m$ be a nonempty S_i -open set. Then

$$\bigcap_{i=1}^n A_i \neq \emptyset.$$

Proof. We shrink each A_i to a product of nonempty S_i -balls $\prod_{j=1}^m B_{S_i}(x_{ij}, z_{ij})$, then for each j we apply Theorem 6.3 in Situation **m** to x_{1j}, \dots, x_{nj} and z_{1j}, \dots, z_{nj} . \square

Theorem 7.7. *Let $S_1, \dots, S_n \subseteq S(K) \setminus \{v_{\text{trivial}}\}$ be nonempty and pairwise disjoint, $\underline{y}_1, \dots, \underline{y}_n \in K^m$ and $\underline{z}_1, \dots, \underline{z}_n \in (K^\times)^m$.*

- (U) *Assume that there exist $f \in K[X]$ and $t \in K^\times$ such that conditions 3.1(i, iii, iv) are satisfied for all $v \in S = S_1 \cup \dots \cup S_n$, and condition 3.1(ii) is satisfied for all $v \in S' = S_1 \cup \dots \cup S_k$, where S_{k+1}, \dots, S_n are singletons.*
- (T) *Assume each S_i is compact in the constructible topology.*
- (I) *Assume that the elements of S are pairwise independent.*

Let $D \subseteq K^l$ which is v -open for all $v \in S$, and let $g : D \rightarrow K^m$ be S' -hereditarily continuous and S_i -continuous for $i = k+1, \dots, n$. Suppose that for each i and $v \in S_i$ there exists some $\underline{x}_v \in D$ with

$$g(\underline{x}_v) \in B_v(\underline{y}_i, \underline{z}_i).$$

Then there exists $\underline{x} \in D$ with

$$g(\underline{x}) \in B_v(\underline{y}_i, \underline{z}_i)$$

for every i and $v \in S_i$.

Proof. For each i and each $v \in S_i$, the condition $g(\underline{x}_v) \in B_v(\underline{y}_i, \underline{z}_i)$ is satisfied in an open-closed neighbourhood of v in the constructible topology on S . By compactness, we find a finite covering of S_i by such open-closed sets. By further refining this covering, we obtain a family $\{S_{i,1}, \dots, S_{i,k_i}\}$ which is a partition of S_i by nonempty open-closed sets, such that for each j there exists $\underline{x}_{ij} \in D$ such that $g(\underline{x}_{ij}) \in B_v(\underline{y}_i, \underline{z}_i)$ for each $v \in S_{i,j}$. Therefore, for each i, j , we have $g(\underline{x}_{ij}) \in B_{S_{i,j}}(\underline{y}_i, \underline{z}_i)$. It follows from our assumptions that $D \subseteq K^l$ is $S_{i,j}$ -open, and likewise it follows that g is $S_{i,j}$ -continuous. Therefore the preimage $A_{i,j} := g^{-1}(B_{S_{i,j}}(\underline{y}_i, \underline{z}_i)) \subseteq D$ is a nonempty $S_{i,j}$ -open set, for each pair i, j . By Proposition 7.6, there exists $x \in \bigcap_{i=1}^n \bigcap_{j=1}^{k_i} A_{i,j}$, and this satisfies the claim. \square

Corollary 7.8. *Let $S_1, \dots, S_n \subseteq S(K) \setminus \{v_{\text{trivial}}\}$ be nonempty and pairwise disjoint, $\underline{y}_1, \dots, \underline{y}_n \in K^m$ and $\underline{z}_1, \dots, \underline{z}_n \in (K^\times)^m$.*

- (U) *Assume that there exist $f \in K[X]$ and $t \in K^\times$ such that conditions 3.1(i, iii, iv) are satisfied for all $v \in S = S_1 \cup \dots \cup S_n$, and condition 3.1(ii) is satisfied for all $v \in S' = S_1 \cup \dots \cup S_k$, where S_{k+1}, \dots, S_n are singletons.*
- (T) *Assume each S_i is compact in the constructible topology.*
- (I) *Assume that the elements of S are pairwise independent.*

Let $g_1, \dots, g_m \in K(x_1, \dots, x_l)$ be rational functions. Suppose that for each i and $v \in S_i$ there exists some $\underline{x}_v \in D_g$ (cf. Example 7.5) with

$$g_j(\underline{x}_v) \in B_v(y_{ij}, z_{ij})$$

for every j . Then there exists $\underline{x} \in D_g$ with

$$g_j(\underline{x}) \in B_v(y_{ij}, z_{ij})$$

for every i, j and $v \in S_i$.

Remark 7.9. We remark that Corollary 7.8 is indeed a generalization of Situation **m** of Theorem 6.3 (so in particular of Theorem 4.6) under the stronger assumption of pairwise independence, as the latter one can be reobtained by taking the g_j to be linear polynomials. We do not know whether Theorem 7.7 or Corollary 7.8 hold without the assumption of independence under some natural compatibility condition.

Remark 7.10. Since in the last theorem and its corollary we have been working with a set S of pairwise independent localities, it is possible to use approximation results from the literature instead of our Theorem 6.3, with a suitable adjustment of the (U) condition. For example, all results of this section will remain valid for sets S_1, \dots, S_n as in Theorem 5.12.

ACKNOWLEDGEMENTS

The authors would like to thank Marcus Tressl for several helpful discussions on spectral spaces and spaces of valuations and orderings.

Part of this work was done while all three authors were guests of the Institut Henri Poincaré and the authors would like to thank the IHP for funding and hospitality, and the organizers of the trimester ‘Model theory, combinatorics and valued fields’ for the invitation.

S. A. was supported by the Leverhulme Trust under grant RPG-2017-179. A. F. was funded by the Deutsche Forschungsgemeinschaft (DFG) - 404427454.

REFERENCES

- [AW45] E. Artin and G. Whaples. Axiomatic characterization of fields by the product formula for valuations. *Bulletin Amer. Math. Soc.* 51 (1945) 469–492.
- [ADF18a] S. Anscombe, P. Dittmann, and A. Fehm. A p -adic analogue of Siegel’s theorem on sums of four squares. Manuscript, 2018.
- [ADF18b] S. Anscombe, P. Dittmann, and A. Fehm. Denseness results in the theory of algebraic fields. Manuscript, 2018.
- [Bou89] N. Bourbaki. *Commutative Algebra. Chapter 1-7*. Springer, 1989.
- [CF67] J. W. S. Cassels and A. Fröhlich. *Algebraic Number Theory*. Academic Press, 1967.
- [Dar98] L. Darnière. *Étude modèle-théorique d’anneaux satisfaisant un principe de Hasse non singulier*. Doctoral thesis, Université de Rennes 1, 1998.
- [Dar00] L. Darnière. Nonsingular Hasse principle for rings. *J. reine angew. Math.* 529:75–100, 2000.
- [DST19] M. Dickmann, N. Schwartz and M. Tressl. *Spectral Spaces*. Cambridge University Press. To appear.
- [Efr06] I. Efrat. *Valuations, Orderings and Milnor K -Theory*. American Mathematical Society, 2006.
- [EP05] A. J. Engler and A. Prestel. *Valued Fields*. Springer, 2005.
- [Ers82] Y. L. Ershov. Totally real field extensions. *Soviet Math. Dokl.*, 25(2):477–480, 1982.
- [Ers94] Y. L. Ershov. Relatively regularly closed fields. *Russian Acad. Sci. Dokl. Math.*, 48(2):300–303, 1994.
- [Ers01] Y. L. Ershov. *Multi-Valued Fields*. Kluwer, 2001.
- [Ers06] Y. L. Ershov. Local-global principles and approximation theorems. In *Logic in Tehran* (eds A. Enayat, I. Kalantari, and M. Moniri), Lecture Notes in Logic, Cambridge University Press, 26:114–125, 2006.
- [Feh13] A. Fehm. Elementary geometric local-global principles for fields. *Annals Pure Appl. Logic*, 164:989–1008, 2013.
- [FHV94] M. Fried, D. Haran and H. Völklein. Real Hilbertianity and the field of totally real numbers. *Contemporary Mathematics* 74:1–34, 1994.
- [GJ91] W. D. Geyer and M. Jarden. The Henselian Closures of a PpC Field. *Abh. Math. Sem. Univ. Hamburg* 61:63–71, 1991.
- [HJP13] D. Haran, M. Jarden and F. Pop. The block approximation theorem. *Math. J. Okayama Univ.* 55(1):53–85, 2013.
- [Hoc69] M. Hochster. Prime ideal structure in commutative rings. *Trans. Amer. Math. Soc.*, 142:43–60, 1969.
- [Kru31] W. Krull. Allgemeine Bewertungstheorie. *J. reine angew. Math.* 167:160–196, 1931.
- [Lam05] T. Y. Lam. *Introduction to Quadratic Forms over Fields*. American Mathematical Society, 2005.
- [Lan02] S. Lang. *Algebra*. Springer, 2002.
- [Mar77] D. A. Marcus. *Number Fields*. Springer, 1977.
- [Nag53] M. Nagata. On the theory of henselian rings. *Nagoya Math. J.* 5:45–57, 1953.
- [Neu99] J. Neukirch. *Algebraic Number Theory*. Springer, 1999.
- [Pre84] A. Prestel. *Formally real fields*. Springer, 1984.
- [Pre85] A. Prestel. On the axiomatization of PRC-fields. In: *Proc. of the VI Latin American Symposium on Mathematical Logic, Caracas 1983*, Springer Lecture Notes in Math. 1130, pp. 351–359, 1985.
- [PR84] A. Prestel and P. Roquette. *Formally p -adic fields*. Springer, 1984.
- [PZ87] A. Prestel and M. Ziegler. Model theoretic methods in the theory of topological fields. *J. reine angew. Math.* 299-300:318–341, 1987.
- [Rib57] P. Ribenboim. Le théorème d’approximation pour les valuations de Krull. *Math. Zeitschr.* 68:1–18, 1957.
- [Rib68] P. Ribenboim. *Theorie des Valuations*. Les Presses de l’Université de Montréal, 1968.
- [Roq73] P. Roquette. Principal ideal theorems for holomorphy rings in fields. *J. reine angew. Math.* 262-263:361–374, 1973.
- [Roq02] P. Roquette. History of Valuation Theory. Part I. In: F.-V. Kuhlmann, S. Kuhlmann, M. Marshall (eds.), *Valuation theory and its applications*, pp. 291–355, 2002.

[War89] S. Warner. *Topological Fields*. North-Holland, 1989.

JEREMIAH HORROCKS INSTITUTE, UNIVERSITY OF CENTRAL LANCASHIRE, PRESTON PR1 2HE, UNITED KINGDOM

E-mail address: sanscombe@uclan.ac.uk

AFDELING ALGEBRA, KU LEUVEN, CELESTIJNENLAAN 200B – BUS 2400, 3001 LEUVEN, BELGIUM

E-mail address: philip.dittmann@kuleuven.be

INSTITUT FÜR ALGEBRA, TECHNISCHE UNIVERSITÄT DRESDEN, 01062 DRESDEN, GERMANY

E-mail address: arno.fehm@tu-dresden.de