

How to determine the *sign* of a valuation on $\mathbb{C}[x, y]$?

Pinaki Mondal
Weizmann Institute of Science
pinaki@math.toronto.edu

December 24, 2013

Abstract

Given a divisorial discrete valuation *centered at infinity* on $\mathbb{C}[x, y]$, we show that its sign on $\mathbb{C}[x, y]$ (i.e. whether it is negative or non-positive on $\mathbb{C}[x, y] \setminus \mathbb{C}$) is completely determined by the sign of its value on the *last key form* (key forms being the avatar of *key polynomials* of valuations [Mac36] in ‘global coordinates’). We also describe the cone of curves and the nef cone of certain compactifications of \mathbb{C}^2 associated to a given valuation centered at infinity, and give a characterization of the divisorial valuations centered at infinity whose *skewness* can be interpreted in terms of the *slope* of an extremal ray of these cones, yielding a generalization of a result of [FJ07]. A by-product of these arguments is a characterization of valuations which ‘determine’ normal compactifications of \mathbb{C}^2 with one irreducible curve at infinity in terms of an associated ‘semigroup of values’.

1 Introduction

Notation 1.1. Throughout this section k is a field and R is a finitely generated k -algebra.

In algebraic (or analytic) geometry and commutative algebra, valuations are usually treated in the *local setting*, and the values are always positive or non-negative. Even if it is a priori not known if a given discrete valuation ν is positive or non-negative on $R \setminus k$, it is evident how to verify this, at least if $\nu(k \setminus \{0\}) = 0$: one has only to check the values of ν on the k -algebra generators of R . For valuations *centered at infinity* however, in general it is non-trivial to determine if it is negative or non-positive on $R \setminus k$:

Example 1.2. Let $R := \mathbb{C}[x, y]$ and for every $\epsilon \in \mathbb{R}$ with $0 < \epsilon < 1$, let ν_ϵ be the valuation (with values in \mathbb{R}) on $\mathbb{C}(x, y)$ defined as follows:

$$\nu_\epsilon(f(x, y)) := -\deg_x \left(f(x, y)|_{y=x^{5/2}+x^{-1}+\xi x^{-5/2-\epsilon}} \right) \quad \text{for all } f \in \mathbb{C}(x, y) \setminus \{0\}, \quad (1)$$

where ξ is a new indeterminate and \deg_x is the degree in x . Direct computation shows that

$$\nu_\epsilon(x) = -1, \quad \nu_\epsilon(y) = -5/2, \quad \nu_\epsilon(y^2 - x^5) = -3/2, \quad \nu_\epsilon(y^2 - x^5 - 2x^{-1}y) = \epsilon.$$

Is ν_ϵ negative on $\mathbb{C}[x, y]$? Let $g := y^2 - x^5 - 2x^{-1}y$. The fact that $\nu_\epsilon(g) > 0$ does not seem to be of much help for the answer (especially if ϵ is very small), since $g \notin \mathbb{C}[x, y]$ and $\nu_\epsilon(xg) < 0$. However, g is precisely the *last key form* (Definition 2.4) of ν_ϵ (see Example 2.8), and therefore Theorem 1.4 implies that ν_ϵ is *not* non-positive on $\mathbb{C}[x, y]$, i.e. no matter how small ϵ is, there exists $f_\epsilon \in \mathbb{C}[x, y]$ such that $\nu_\epsilon(f_\epsilon) > 0$.

In this article we settle the question of how to determine if a valuation centered at infinity is negative or non-positive on R for the case that $R = \mathbb{C}[x, y]$. At first we describe how this question arises naturally in the study of algebraic completions of affine varieties.

1.1 Motivation and statements of main results

Recall that a *divisorial discrete valuation* (Definition 2.2) ν on R is *centered at infinity* iff $\nu(f) < 0$ for some $f \in R$, or equivalently iff there is an *algebraic completion* \bar{X} of $X := \text{Spec } R$ (i.e. \bar{X} is a complete algebraic varieties containing X as a dense open subset) and an irreducible component C of $\bar{X} \setminus X$ such that ν is the order of vanishing along C . On the other hand, one way to *construct* algebraic completions of the affine variety X is to start with a *degree-like function* on R (the terminology is from [Mon10b] and [Mon10a]), i.e. a function $\delta : R \rightarrow \mathbb{Z} \cup \{-\infty\}$ which satisfy the following ‘degree-like’ properties:

P1. $\delta(f + g) \leq \max\{\delta(f), \delta(g)\}$, and

P2. $\delta(fg) \leq \delta(f) + \delta(g)$,

and construct the graded ring

$$R^\delta := \bigoplus_{d \geq 0} \{f \in R : \delta(f) \leq d\} \cong \sum_{d \geq 0} \{f \in R : \delta(f) \leq d\} t^d \quad (2)$$

where t is an indeterminate. It is straightforward to see that $\bar{X}^\delta := \text{Proj } R^\delta$ is a *projective completion* of X provided the following conditions are satisfied:

Proj-1. R^δ is finitely generated as a k -algebra, and

Proj-2. $\delta(f) > 0$ for all $f \in R \setminus k$.

A fundamental class of degree-like functions are *divisorial semidegrees* - these are precisely the *negative* of divisorial discrete valuations centered at infinity and they serve as ‘building blocks’ of an important class of degree-like functions (see [Mon10b], [Mon10a]). Therefore, a natural question in this context is:

Question 1.3. ¹ Given a divisorial semidegree δ on R , how to determine if $\delta(f) > 0$ for all $f \in R \setminus k$? Or equivalently, given a divisorial discrete valuation ν on R centered at infinity, how to determine if $\nu(f) < 0$ for all $f \in R \setminus k$?

In this article we give a complete answer to Question 1.3 for the case $k = \mathbb{C}$ and $R = \mathbb{C}[x, y]$ (note that the answer for the case $R = \mathbb{C}[x]$ is obvious, since the only discrete valuations centered at infinity on $\mathbb{C}[x]$ are those which map $x - \alpha \mapsto -1$ for some $\alpha \in \mathbb{C}$). More precisely, we consider the sequence of *key forms* (Definition 2.4) corresponding to semidegrees, and show that

Theorem 1.4. *Let δ be a **divisorial semidegree** on $\mathbb{C}[x, y]$ and let g_0, \dots, g_{n+1} be the key forms of δ in (x, y) -coordinates. Then*

¹The analogous question regarding Property (Proj-1) for $R = \mathbb{C}[x, y]$ and δ a divisorial semidegree is completely settled in [MN13] where the results of this article are also applied to the *moment problem* of planar semialgebraic sets.

1. δ is non-negative on $\mathbb{C}[x, y] \setminus \mathbb{C}$ iff $\delta(g_{n+1})$ is non-negative.
2. δ is positive on $\mathbb{C}[x, y] \setminus \mathbb{C}$ iff one of the following holds:
 - (a) $\delta(g_{n+1})$ is positive,
 - (b) $\delta(g_{n+1}) = 0$ and $g_k \notin \mathbb{C}[x, y]$ for some k , $0 \leq k \leq n + 1$, or
 - (b') $\delta(g_{n+1}) = 0$ and $g_{n+1} \notin \mathbb{C}[x, y]$.

Moreover, conditions **2b** and **2b'** are equivalent.

Remark 1.5. The key forms of a **semidegree** δ on $\mathbb{C}[x, y]$ are counterparts in (x, y) -coordinates of the *key polynomials* of $\nu := -\delta$ introduced in [Mac36] (and computed in local coordinates near the *center* of ν). The basic ingredient of the proof of Theorem 1.4 is the algebraic contraitibility criterion of [Mon13a] which uses key forms. We note that key forms were already used in [FJ07]² (without calling them by any special name).

Remark 1.6. The requirement in Theorem 1.4 that δ be *divisorial* (i.e. $-\delta$ be a *divisorial* valuation) is unnecessary: the only technical issue stems from valuations with an infinite sequence of key polynomials - but one can determine the sign of such a valuation by applying Theorem 1.4 to a divisorial valuation which ‘approximates’ it sufficiently closely.

Remark 1.7. The key forms of a **semidegree** can be computed explicitly from any of the alternative presentations of the semidegree (see e.g. [Mon13a, Algorithm 3.24] for an algorithm to compute key forms from the *generic Puiseux series* (Definition 2.13) associated to the semidegree). Therefore Theorem 1.4 gives an *effective* way to determine if a given semidegree is positive or non-negative on $\mathbb{C}[x, y]$.

Trees of valuations centered at infinity on $\mathbb{C}[x, y]$ were considered in [FJ07] along with a parametrization of the tree called *skewness* α . The notion of skewness has an ‘obvious’ extension³ to the case of semidegrees, and using this definition one of the assertions of [FJ07, Theorem A.7] can be reformulated as the statement that the following identity holds for a certain *subtree* of semidegrees δ on $\mathbb{C}[x, y]$:

$$\alpha(\delta) = \inf \left\{ \frac{\delta(f)}{d_\delta \deg(f)} : f \text{ is a non-constant polynomial in } \mathbb{C}[x, y] \right\}, \text{ where} \quad (3)$$

$$d_\delta := \max\{\delta(x), \delta(y)\}. \quad (4)$$

It is observed in [Jon12, Page 121] that in general the relation in (3) is satisfied with \leq , and “it is doubtful that equality holds in general.” Example 3.1 shows that the equality indeed does not hold in general. It is not hard to see that $\alpha(\delta)$ can be expressed in terms of $\delta(g_{n+1})$ (see (17)), and using that expression we give a characterization of the semidegrees for which (3) holds true:

Theorem 1.8. *Let δ be a **semidegree** on $\mathbb{C}[x, y]$ and g_0, \dots, g_{n+1} be the corresponding key forms. Then (3) holds iff one of the following assertions is true:*

²Under the assumptions of Lemma A.12 of [FJ07], the polynomials U_j constructed in Section A.5.3 of [FJ07] are precisely the key forms of $-\nu$.

³In [FJ07] the skewness α was defined only for valuations ν centered at infinity which satisfied $\min\{\nu(x), \nu(y)\} = -1$. Here for a semidegree δ , we define $\alpha(\delta)$ to be the skewness of $-\delta/d_\delta$ (where d_δ is as in (4)) in the sense of [FJ07].

1. $\delta(g_{n+1}) \geq 0$, or
2. $\delta(g_{n+1}) < 0$ and $g_k \in \mathbb{C}[x, y]$ for all k , $0 \leq k \leq n + 1$, or
- 2'. $\delta(g_{n+1}) < 0$ and $g_{n+1} \in \mathbb{C}[x, y]$.

Moreover, the ‘inf’ in right hand side of (3) can be replaced by ‘min’ iff $g_{n+1} \in \mathbb{C}[x, y]$ iff $g_k \in \mathbb{C}[x, y]$ for all k , $0 \leq k \leq n + 1$; in this case the minimum is achieved with $f = g_{n+1}$.

Remark 1.9. It is possible to give a *geometric* characterization of the semidegrees δ for which (3) holds. Indeed, [CPR05] introduced the notion of compactifications of \mathbb{C}^2 which admit systems of numerical curvettes. In Section 1.2 and Remark 1.13 below we construct two compactifications \bar{X} and \tilde{X} of \mathbb{C}^2 associated to δ . [Mon13b, Theorem 3.2] (which uses the results of this article), shows that (3) holds iff \bar{X} (or equivalently, \tilde{X}) admits a system of numerical curvettes.

Our final result is the following corollary of the arguments in the proof of Theorem 1.4 which answers a question of Professor Peter Russell⁴.

Corollary 1.10. *Let δ be a semidegree on $\mathbb{C}[x, y]$. Define*

$$S_\delta := \{(\deg(f), \delta(f)) : f \in \mathbb{C}[x, y] \setminus \{0\}\} \subseteq \mathbb{Z}^2, \quad (5)$$

and \mathcal{C}_δ be the cone over S_δ in \mathbb{R}^2 . Then

1. The following are equivalent:
 - (a) \mathcal{C}_δ is a closed subset of \mathbb{R}^2 .
 - (b) g_k is a polynomial for all k , $0 \leq k \leq n + 1$.
 - (c) g_{n+1} is a polynomial.
2. The following are equivalent:
 - (a) δ determines an analytic compactification of \mathbb{C}^2 .
 - (b) the positive x -axis is not contained in the closure $\bar{\mathcal{C}}_\delta$ of \mathcal{C}_δ in \mathbb{R}^2 .
3. The following are equivalent:
 - (a) δ determines an algebraic compactification of \mathbb{C}^2 .
 - (b) \mathcal{C}_δ is closed in \mathbb{R}^2 and the positive x -axis is not contained in \mathcal{C}_δ .
 - (c) S_δ is a finitely generated semigroup and $(k, 0) \notin S_\delta$ for all positive integer k .

⁴Prof. Russell’s question was motivated by the correspondence established in [Mon13a] between normal algebraic compactifications of \mathbb{C}^2 with one irreducible curve at infinity and algebraic curves in \mathbb{C}^2 with one place at infinity. Since the *semigroup of poles* of planar curves with one place at infinity are very *special* (see e.g. [Abh78], [SS94]), he asked if similarly the semigroups of values of semidegrees which determine normal algebraic compactifications of \mathbb{C}^2 can be similarly distinguished from the semigroup of values of general semidegrees. While Example 3.2 shows that they can not be distinguished only by the values of the semidegree itself, Corollary 1.10 shows that it can be done if paired with degree of polynomials.

Remark 1.11. The phrase “ δ determines an algebraic (resp. analytic) compactification of \mathbb{C}^2 ” means “there exists a (necessarily unique) normal algebraic (resp. analytic) compactification \bar{X} of $X := \mathbb{C}^2$ such that $C_\infty := \bar{X} \setminus X$ is an irreducible curve and δ is proportional to the order of pole along C_∞ .” In particular, δ determines an algebraic compactification of \mathbb{C}^2 iff δ satisfies conditions [Proj-1](#) and [Proj-2](#).

Remark 1.12. S_δ is isomorphic to the *global Enriques semigroup* (in the terminology of [\[CPRL02\]](#)) of the compactification of \mathbb{C}^2 from [Proposition 2.10](#). Also, the assertions of [Corollary 1.10](#) remain true if in [\(5\)](#) deg is replaced by any other semidegree which determines an algebraic completion of \mathbb{C}^2 (e.g. a weighted degree with positive weights).

1.2 Cones of curves on compactifications of \mathbb{C}^2

Let $\mathbb{C}^2 \subseteq \mathbb{P}^2 = \mathbb{C}^2 \cup L$ be the usual compactification of \mathbb{C}^2 (where L is the ‘line at infinity’) and δ be a **divisorial semidegree** on $\mathbb{C}[x, y]$ centered at infinity. Assume $\delta \neq \text{deg}$. By local theory, there exists a birational map $\pi : \bar{X} \rightarrow \mathbb{P}^2$, \bar{X} normal \mathbb{Q} -factorial, such that the center of δ on \bar{X} is the whole reduced (irreducible) exceptional curve C_2 ([Proposition 2.10](#)). Let C_1 be the strict transform of L on \bar{X} . For a curve $D \subseteq \mathbb{C}^2$ defined by a polynomial $f \in \mathbb{C}[x, y]$, its closure \bar{D} in \bar{X} is linearly equivalent to $\text{deg}(f)C_1 + \delta(f)C_2$ as Weil divisors on \bar{X} . It follows that the group $N_1(\bar{X})$ of Weil divisors on \bar{X} modulo numerical equivalence is a free group generated by C_1 and C_2 . [Theorem 1.4](#) is related to the question of whether C_1 and C_2 generate the cone $\text{NE}(\bar{X})$ of curves on \bar{X} . More precisely, an ingredient of [Theorem 1.4](#) is the following result:

Theorem 1.4’. *Let g_0, \dots, g_{n+1} be the key forms of δ in (x, y) -coordinates. Then*

1. $\delta(g_{n+1})$ equals a negative rational number times the self intersection number of C_1 .
2. $\delta(g_{n+1}) \geq 0$ iff C_1 and C_2 generate $\text{NE}(\bar{X})$. Let l_i be the half-line of all non-negative real multiples of C_i , $1 \leq i \leq 2$. Then,
 - (a) l_2 determines an edge of $\text{NE}(\bar{X})$.
 - (b) l_1 determines an edge of $\text{NE}(\bar{X})$ iff $\delta(g_{n+1}) \geq 0$.
 - (c) C_1 is in the interior of $\text{NE}(\bar{X})$ iff $\delta(g_{n+1}) < 0$.

Similarly, [Theorem 1.8](#) is related to properties of the nef cone $\text{Nef}(\bar{X})$ of \bar{X} . More precisely, let g_0, \dots, g_{n+1} be the key forms of δ in (x, y) -coordinates. Define \mathbb{Q} -Cartier divisors $C_1^* := C_1 + d_\delta C_2$ and $C_2^* := C_1 + \frac{m_\delta \delta(g_{n+1})}{d_\delta} C_2$ on \bar{X} , where d_δ is as in [\(4\)](#), and

$$m_\delta := \gcd(\delta(g_0), \dots, \delta(g_n)). \quad (6)$$

Let l_i^* be the half-line of all non-negative real multiples of C_i^* , $1 \leq i \leq 2$.

Theorem 1.8’. $\delta(g_{n+1}) \geq 0$ iff C_1^* and C_2^* generate $\text{Nef}(\bar{X})$. More precisely

1. l_1^* determines an edge of $\text{Nef}(\bar{X})$.
2. l_2^* determines an edge of $\text{Nef}(\bar{X})$ iff $\delta(g_{n+1}) \geq 0$.
3. $C_2^* \notin \text{Nef}(\bar{X})$ iff $\delta(g_{n+1}) < 0$.

Remark 1.13. Consider the minimal resolution of singularities $\tilde{\pi} : \tilde{X} \rightarrow \bar{X}$. Let \tilde{E} be the union of the exceptional curves of $\tilde{\pi}$ with the strict transform of C_1 . Then assertion 1 of Theorem 1.4' implies that

$$\begin{aligned} \delta(g_{n+1}) > 0 & \text{ iff } (C_1, C_1) < 0 \\ & \text{ iff the intersection matrix of } \tilde{E} \text{ is negative definite,} \end{aligned} \tag{7}$$

$$\begin{aligned} \delta(g_{n+1}) \geq 0 & \text{ iff } (C_1, C_1) \leq 0 \\ & \text{ iff the intersection matrix of } \tilde{E} \text{ is nonpositive definite.} \end{aligned} \tag{8}$$

In particular, the property that $\delta(g_{n+1}) > 0$ (resp. $\delta(g_{n+1}) \geq 0$) is equivalent to a purely numerical criterion (7) (resp. (8)) which is completely determined by the weighted configuration of projective lines on $\tilde{X} \setminus \mathbb{C}^2$.

Remark 1.14. Let g_0, \dots, g_{n+1} be the key forms of δ in (x, y) -coordinates, and let \tilde{X} be as in Remark 1.13. Let $\tilde{C}_1, \dots, \tilde{C}_k$ be the irreducible components of $\tilde{X} \setminus \mathbb{C}^2$ and for each j , $0 \leq j \leq k$, let δ_j be the semidegree on $\mathbb{C}[x, y]$ associated to \tilde{C}_j . It is not hard to see that the last key form of δ_j is g_{i_j} for some i_j , $1 \leq i_j \leq n+1$. Moreover, in the case that $\delta(g_{n+1}) \geq 0$, it turns out that $\delta_j(g_{i_j}) \geq 0$ for each j , $1 \leq j \leq k$. Theorem 1.4 then implies that $\text{NE}(\tilde{X})$ is (the simplicial cone) generated by $\tilde{C}_1, \dots, \tilde{C}_k$. Combining this with Remark 1.13 it follows that if $\delta(g_{n+1}) \geq 0$ then the cones $\text{NE}(\tilde{X})$ and $\text{Nef}(\tilde{X})$ are (simplicial and) completely determined by the weighted configuration of projective lines on $\tilde{X} \setminus \mathbb{C}^2$. However, if $\delta(g_{n+1}) < 0$, Example 3.3 below shows that in general the weighted configuration of projective lines on $\tilde{X} \setminus \mathbb{C}^2$ does not determine $\text{NE}(\tilde{X})$ or $\text{Nef}(\tilde{X})$.

1.3 Acknowledgements

I am grateful to Professor Peter Russell for the question which led to this work. I would like to thank Professor Pierre Milman - the mathematics of this article was worked out while I was his postdoc at University of Toronto. I would also like to thank the referee for his/her valuable suggestions which contributed significantly to the clarity of the underlying geometry. The article has been written up at the Weizmann Institute as an Azrieli Fellow. I am grateful to the Azrieli Foundation for the award of an Azrieli Fellowship.

2 Preliminaries

Notation 2.1. Throughout the rest of the article we write $X := \mathbb{C}^2$ with polynomial coordinates (x, y) and let $\bar{X}^{(0)} \cong \mathbb{P}^2$ be the compactification of X induced by the embedding $(x, y) \mapsto [1 : x : y]$, so that the semidegree on $\mathbb{C}[x, y]$ corresponding to the line at infinity is precisely on $\bar{X}^{(0)}$ is deg , where deg is the usual degree in (x, y) -coordinates.

2.1 Divisorial discrete valuations, semidegrees, key forms, and associated compactifications

Definition 2.2 (Discrete valuations). A discrete valuation on $\mathbb{C}(x, y)$ is a map $\nu : \mathbb{C}(x, y) \setminus \{0\} \rightarrow \mathbb{Z}$ such that for all $f, g \in \mathbb{C}(x, y) \setminus \{0\}$,

1. $\nu(f + g) \geq \min\{\nu(f), \nu(g)\}$,
2. $\nu(fg) = \nu(f) + \nu(g)$.

A discrete valuation ν on $\mathbb{C}(x, y)$ is called *divisorial* iff there exists a normal algebraic surface Y_ν equipped with a birational map $\sigma : Y_\nu \rightarrow \bar{X}^0$ and a curve C_ν on Y_ν such that for all non-zero $f \in \mathbb{C}[x, y]$, $\nu(f)$ is the order of vanishing of $\sigma^*(f)$ along C_ν . The *center* of ν on \bar{X}^0 is $\sigma(C_\nu)$. ν is said to be *centered at infinity* (with respect to (x, y) -coordinates) iff the center of ν on \bar{X}^0 is contained in $\bar{X}^0 \setminus X$; equivalently, ν is centered at infinity iff there is a non-zero polynomial $f \in \mathbb{C}[x, y]$ such that $\nu(f) < 0$.

Definition 2.3 (Semidegrees). A (*divisorial*) *semidegree* on $\mathbb{C}(x, y) \setminus \{0\}$ is a map $\delta : \mathbb{C}(x, y) \setminus \{0\} \rightarrow \mathbb{Z}$ such that $-\delta$ is a (divisorial) discrete valuation centered at infinity.

Definition 2.4 (cf. definition of key polynomials in [FJ04, Definition 2.1], also see Remark 2.6 below). Let δ be a divisorial semidegree on $\mathbb{C}[x, y]$ such that $\delta(x) > 0$. A sequence of elements $g_0, g_1, \dots, g_{n+1} \in \mathbb{C}[x, x^{-1}, y]$ is called the sequence of *key forms* for δ if the following properties are satisfied:

P0. $g_0 = x, g_1 = y$.

P1. Let $\omega_j := \delta(g_j)$, $0 \leq j \leq n+1$. Then

$$\omega_{j+1} < \alpha_j \omega_j = \sum_{i=0}^{j-1} \beta_{j,i} \omega_i \text{ for } 1 \leq j \leq n,$$

where

(a) $\alpha_j = \min\{\alpha \in \mathbb{Z}_{>0} : \alpha \omega_j \in \mathbb{Z}\omega_0 + \dots + \mathbb{Z}\omega_{j-1}\}$ for $1 \leq j \leq n$,

(b) $\beta_{j,i}$'s are integers such that $0 \leq \beta_{j,i} < \alpha_j$ for $1 \leq i < j \leq n$ (in particular, $\beta_{j,0}$'s are allowed to be *negative*).

P2. For $1 \leq j \leq n$, there exists $\theta_j \in \mathbb{C}^*$ such that

$$g_{j+1} = g_j^{\alpha_j} - \theta_j g_0^{\beta_{j,0}} \dots g_{j-1}^{\beta_{j,j-1}}.$$

P3. Let y_1, \dots, y_{n+1} be indeterminates and ω be the *weighted degree* on $B := \mathbb{C}[x, x^{-1}, y_1, \dots, y_{n+1}]$ corresponding to weights ω_0 for x and ω_j for y_j , $0 \leq j \leq n+1$ (i.e. the value of ω on a polynomial is the maximum ‘weight’ of its monomials). Then for every polynomial $g \in \mathbb{C}[x, x^{-1}, y]$,

$$\delta(g) = \min\{\omega(G) : G(x, y_1, \dots, y_{n+1}) \in B, G(x, g_1, \dots, g_{n+1}) = g\}. \quad (9)$$

Theorem 2.5 ([Mon13a, Theorem 3.18], cf. [FJ04, Theorem 2.29]). *There is a unique and finite sequence of key forms for δ .*

Remark 2.6. Let δ be as in Definition 2.4. Set $u := 1/x$ and $v := y/x^k$ for some k such that $\delta(y) < k\delta(x)$, and let $\tilde{g}_0 = u, \tilde{g}_1 = v, \tilde{g}_2, \dots, \tilde{g}_{n+1} \in \mathbb{C}[u, v]$ be the *key polynomials* of $\nu := -\delta$ in (u, v) -coordinates. Then the key forms of δ can be computed from \tilde{g}_j 's as follows:

$$g_j(x, y) := \begin{cases} x & \text{for } j = 0, \\ x^{k \deg_v(\tilde{g}_j)} \tilde{g}_j(1/x, y/x^k) & \text{for } 1 \leq j \leq n+1. \end{cases} \quad (10)$$

Theorem 2.5 is an immediate consequence of the existence of key polynomials (see e.g. [FJ04, Theorem 2.29]).

Example 2.7. Let (p, q) are integers such that $p > 0$ and δ be the weighted degree on $\mathbb{C}(x, y)$ corresponding to weights p for x and q for y . Then the key forms of δ are x, y .

Example 2.8. Let $\epsilon := q/2p$ for positive integers p, q such that $q < 2p$ and δ_ϵ be the semidegree on $\mathbb{C}(x, y)$ defined as follows:

$$\delta_\epsilon(f(x, y)) := 2p \deg_x \left(f(x, y)|_{y=x^{5/2}+x^{-1}+\xi x^{-5/2-\epsilon}} \right) \quad \text{for all } f \in \mathbb{C}(x, y) \setminus \{0\}, \quad (11)$$

where ξ is a new indeterminate and \deg_x is the degree in x . Note that $\delta_\epsilon = -2p\nu_\epsilon$, where ν_ϵ is from Example 1.2 (we multiplied by $2p$ to simply make the semidegree integer valued). Then the sequence of key forms of δ_ϵ is $x, y, y^2 - x^5, y^2 - x^5 - 2x^{-1}y$.

The following property of key forms can be proved in a straightforward way from their defining properties.

Proposition 2.9. *Let δ and g_0, \dots, g_{n+1} be as in Definition 2.4, and d_δ and m_δ be as in respectively (4) and (6). Then*

$$m_\delta \delta(g_{n+1}) \leq d_\delta^2. \quad (12)$$

Moreover, (12) is satisfied with an equality iff $\delta = \deg$.

Let $\bar{X}^0 := \mathbb{P}^2$ be the usual compactification of \mathbb{C}^2 given by $(x, y) \mapsto [1 : x : y]$. If δ is a divisorial valuation δ on $\mathbb{C}[x, y]$, then by definition there is a compactification \bar{X}^1 of \mathbb{C}^2 such that δ is the order of pole along an irreducible curve $C \subseteq \bar{X}^1 \setminus \mathbb{C}^2$. W.l.o.g. we may assume that \bar{X}^1 is non-singular and there is a morphism $\pi : \bar{X}^1 \rightarrow \bar{X}^0$ which is identity on \mathbb{C}^2 . In particular, C is an *exceptional curve* of π (i.e. $\pi(C)$ is a point). Let \bar{X} be the surface obtained from \bar{X}^1 by contracting all exceptional curves of π other than C (which is possible due to a criterion of Grauert [Băd01, Theorem 14.20]). Then $\bar{X} \setminus \mathbb{C}^2$ is the union of two irreducible curves, and the following result, which follows from results of [Mon11], describes the matrix of intersection numbers of these curves in terms of the key forms of δ .

Proposition 2.10 ([Mon11, Propositions 4.2 and 4.7]). *Given a divisorial semidegree δ on $\mathbb{C}[x, y]$ such that $\delta \neq \deg$ and $\delta(x) > 0$, there exists a unique compactification \bar{X} of \mathbb{C}^2 such that*

1. \bar{X} is projective and normal.
2. $\bar{X}_\infty := \bar{X} \setminus X$ has two irreducible components C_1, C_2 .
3. The semidegree on $\mathbb{C}[x, y]$ corresponding to C_1 and C_2 are respectively \deg and δ .

Moreover, all singularities \bar{X} are rational (which implies in particular that all Weil divisors are \mathbb{Q} -Cartier). Let g_0, \dots, g_{n+1} be the key forms of δ . Then the inverse of the matrix of intersection numbers (C_i, C_j) of C_i and C_j , $1 \leq i, j \leq 2$, is

$$\mathcal{M} = \begin{pmatrix} 1 & d_\delta \\ d_\delta & m_\delta \delta(g_{n+1}) \end{pmatrix}, \quad (13)$$

where d_δ and m_δ are as in respectively (4) and (6).

We will use the following result which is an immediate corollary of [Mon13a, Proposition 4.2].

Proposition 2.11. *Let δ , \bar{X} and C_1, C_2 be as in Proposition 2.10. Let g_0, \dots, g_{n+1} be the key forms of δ . Then the following are equivalent:*

1. *there is a (compact algebraic) curve C on \bar{X} such that $C \cap C_1 = \emptyset$.*
2. *g_k is a polynomial for all k , $0 \leq k \leq n+1$.*
3. *g_{n+1} is a polynomial.*

The following is the main result of [Mon13a]:

Theorem 2.12. *Let δ be a divisorial semidegree on $\mathbb{C}[x, y]$ such that $\delta(x) > 0$ and g_0, \dots, g_{n+1} be the key forms of δ . Then δ determines a normal algebraic compactification of \mathbb{C}^2 (in the sense of Remark 1.11) iff $\delta(g_{n+1}) > 0$ and g_{n+1} is a polynomial.*

2.2 Degree-wise Puiseux series

Note. The proofs of Theorems 1.4, 1.4' and 1.8' do *not* use the material of this subsection. Proposition 2.20 and Corollary 2.22 are used in the proof of $\delta(g_{n+1}) < 0$ case of Theorem 1.8.

Definition 2.13 (Degree-wise Puiseux series). The field of *degree-wise Puiseux series* in x is

$$\mathbb{C}\langle\langle x \rangle\rangle := \bigcup_{p=1}^{\infty} \mathbb{C}((x^{-1/p})) = \left\{ \sum_{j \leq k} a_j x^{j/p} : k, p \in \mathbb{Z}, p \geq 1 \right\},$$

where for each integer $p \geq 1$, $\mathbb{C}((x^{-1/p}))$ denotes the field of Laurent series in $x^{-1/p}$. Let $\phi = \sum_{q \leq q_0} a_q x^{q/p}$ be a degree-wise Puiseux series where p is the *polydromy order* of ϕ , i.e. p is the smallest positive integer such that $\phi \in \mathbb{C}((x^{-1/p}))$. Then the *conjugates* of ϕ are $\phi_j := \sum_{q \leq q_0} a_q \zeta^q x^{q/p}$, $1 \leq j \leq p$, where ζ is a primitive p -th root of unity. The usual factorization of polynomials in terms of Puiseux series implies the following

Theorem 2.14. *Let $f \in \mathbb{C}[x, y]$. Then there are unique (up to conjugacy) degree-wise Puiseux series ϕ_1, \dots, ϕ_k , a unique non-negative integer m and $c \in \mathbb{C}^*$ such that*

$$f = cx^m \prod_{i=1}^k \prod_{\phi_{ij} \text{ is a conjugate of } \phi_i} (y - \phi_{ij}(x))$$

The relation between degree-wise Puiseux series and semidegrees is given by the following proposition, which is a reformulation of the corresponding result for Puiseux series and valuations [FJ04, Proposition 4.1].

Proposition 2.15 ([Mon11, Theorem 1.2]). *Let δ be a divisorial semidegree on $\mathbb{C}(x, y)$ such that $\delta(x) > 0$. Then there exists a degree-wise Puiseux polynomial (i.e. a degree-wise Puiseux series with finitely many terms) $\phi_\delta \in \mathbb{C}\langle\langle x \rangle\rangle$ and a rational number $r_\delta < \text{ord}_x(\phi_\delta)$ such that for every polynomial $f \in \mathbb{C}[x, y]$,*

$$\delta(f) = \delta(x) \deg_x (f(x, y)|_{y=\phi_\delta(x)+\xi x^{r_\delta}}), \quad (14)$$

where ξ is an indeterminate.

Definition 2.16. If ϕ_δ and r_δ are as in Proposition 2.15, we say that $\tilde{\phi}_\delta(x, \xi) := \phi_\delta(x) + \xi x^{r_\delta}$ is the *generic degree-wise Puiseux series* associated to δ .

Example 2.17. Let (p, q) are integers such that $p > 0$ and δ be the weighted degree on $\mathbb{C}(x, y)$ corresponding to weights p for x and q for y . Then $\tilde{\phi}_\delta = \xi x^{q/p}$ (i.e. $\phi_\delta = 0$).

Example 2.18. Let δ_ϵ be the semidegree from Example 2.8. Then $\tilde{\phi}_\delta = x^{5/2} + x^{-1} + \xi x^{-5/2}$.

The following result, which is an immediate consequence of [Mon11, Proposition 4.2, Assertion 2], connects degree-wise Puiseux series of a semidegree with the geometry of associated compactifications.

Proposition 2.19. Let $\delta, \bar{X}, C_1, C_2$ be as in Proposition 2.10 and let $\tilde{\phi}_\delta(x, \xi) := \phi_\delta(x) + \xi x^{r_\delta}$ be the generic degree-wise Puiseux series associated to δ . Assume in addition that δ is not a weighted degree, i.e. $\phi_\delta(x) \neq 0$. Pick $f \in \mathbb{C}[x, y] \setminus \{0\}$ and let C_f be the curve on \bar{X} which is the closure of the curve defined by f on \mathbb{C}^2 . Then $C_f \cap C_1 = \emptyset$ iff the degree-wise Puiseux factorization of f is of the form

$$f = \prod_{i=1}^k \prod_{\phi_{ij} \text{ is a conjugate of } \phi_i} (y - \phi_{ij}(x)), \quad \text{where each } \phi_i \text{ satisfies} \quad (15)$$

$$\phi_i(x) - \phi_\delta(x) = c_i x^{r_\delta} + \text{l.o.t.}$$

for some $c_i \in \mathbb{C}$ (where l.o.t. denotes lower order terms in x).

The following result gives some relations between degree-wise Puiseux series and key forms of semidegrees, and follows from standard properties of key polynomials (in particular, the first 3 assertions follow from [Mon13a, Proposition 3.28] and the last assertion follows from the first; a special case of the last assertion (namely the case that $\delta(y) \leq \delta(x)$) was proved in [Mon11, Identity (4.6)]).

Proposition 2.20. Let δ be a divisorial semidegree on $\mathbb{C}(x, y)$ such that $\delta(x) > 0$. Let $\tilde{\phi}_\delta(x, \xi) := \phi_\delta(x) + \xi x^{r_\delta}$ be the generic degree-wise Puiseux series associated to δ and g_0, \dots, g_{n+1} be the key forms of δ . Then

1. There is a degree-wise Puiseux series ϕ with

$$\phi(x) - \phi_\delta(x) = c x^{r_\delta} + \text{l.o.t.}$$

for some $c \in \mathbb{C}$ (where l.o.t. denotes lower order terms in x) such that the degree-wise Puiseux factorization of g_{n+1} is of the form

$$g_{n+1} = \prod_{\phi^* \text{ is a conjugate of } \phi} (y - \phi^*(x)). \quad (16)$$

2. Let the Puiseux pairs [Mon13a, Definition 3.11] of ϕ_δ be $(q_1, p_1), \dots, (q_l, p_l)$ (if $\phi_\delta \in \mathbb{C}((1/x))$, then simply set $l = 0$). Set $p_0 := 1$. Then

$$\deg(g_{n+1}) = \begin{cases} 1 & \text{if } \phi_\delta = 0, \\ \max\{1, \deg_x(\phi_\delta)\} p_0 p_1 \cdots p_l & \text{otherwise.} \end{cases}$$

3. Write r_δ as $r_\delta = q_{l+1}/(p_0 \cdots p_l p_{l+1})$, where p_{l+1} is the smallest integer ≥ 1 such that $p_0 \cdots p_l p_{l+1} r_\delta$ is an integer. Let d_δ and m_δ be as in respectively (4) and (6). Then

$$m_\delta = p_{l+1},$$

$$d_\delta = \begin{cases} \max\{p_1, q_1\} & \text{if } \phi_\delta = 0, \\ \max\{1, \deg_x(\phi_\delta)\} p_0 p_1 \cdots p_{l+1} & \text{otherwise.} \end{cases}$$

4. Let the skewness $\alpha(\delta)$ of δ be defined as in footnote 3. Then

$$\alpha(\delta) = m_\delta \delta(g_{n+1}) / d_\delta^2 = \begin{cases} \frac{\min\{p_1, q_1\}}{\max\{p_1, q_1\}} = \min\{\delta(x), \delta(y)\} / d_\delta & \text{if } \phi_\delta = 0, \\ \frac{\delta(g_{n+1})}{d_\delta \deg(g_{n+1})} & \text{otherwise.} \end{cases} \quad (17)$$

The following lemma is a consequence of Assertion 1 of Proposition 2.20 and the definition of generic degree-wise Puiseux series of a semidegree. It follows via a straightforward, but cumbersome induction on the number of *Puiseux pairs* of the *degree-wise Puiseux roots* of f , and we omit the proof.

Lemma 2.21. *Let δ be a divisorial semidegree on $\mathbb{C}(x, y)$ such that $\delta(x) > 0$. Let $\tilde{\phi}_\delta(x, \xi) := \phi_\delta(x) + \xi x^{r_\delta}$ be the generic degree-wise Puiseux series associated to δ and g_0, \dots, g_{n+1} be the key forms of δ . Then for all $f \in \mathbb{C}[x, y] \setminus \mathbb{C}$,*

$$\frac{\delta(f)}{\deg(f)} \geq \frac{\delta(g_{n+1})}{\deg(g_{n+1})}. \quad (18)$$

Now assume in addition that δ is not a weighted degree, i.e. $\phi_\delta(x) \neq 0$. Then (18) holds with equality iff f has a degree-wise Puiseux factorization as in (15).

Combining Propositions 2.11 and 2.19 and Lemma 2.21 yields the following

Corollary 2.22. *Consider the set-up of Proposition 2.11. assume in addition that δ is not a weighted degree. Then the Assertions 1 to 3 of Proposition 2.11 are equivalent to the following statement*

4. *There exists $f \in \mathbb{C}[x, y] \setminus \mathbb{C}$ which satisfies (18) with equality.*

3 Proofs

Proof of Theorem 1.4'. Let \bar{X} be the projective compactification of $X := \mathbb{C}^2$ from Section 1.2. In the notations of Proposition 2.10, the matrix of intersection numbers (C_i, C_j) of C_i and C_j , $1 \leq i, j \leq 2$, is:

$$\mathcal{I} = \frac{1}{d_\delta^2 - m_\delta \delta(g_{n+1})} \begin{pmatrix} -m_\delta \delta(g_{n+1}) & d_\delta \\ d_\delta & -1 \end{pmatrix} \quad (19)$$

In particular, $(C_1, C_1) = -\frac{m_\delta}{d_\delta^2 - m_\delta \delta(g_{n+1})} \delta(g_{n+1})$. Since $\delta \neq \deg$ (by the assumptions of Theorem 1.4'), assertion 1 of Theorem 1.4' follows from Proposition 2.9. It follows similarly that

$(C_2, C_2) < 0$, so that [Kol96, Lemma II.4.12]⁵ implies that l_2 determines an edge of $\text{NE}(\bar{X})$, which implies assertion 2a. Now observe that

$$\begin{aligned} \delta(g_{n+1}) \geq 0 &\Rightarrow (C_1, C_1) \leq 0 \text{ (assertion 1)} \\ &\Rightarrow l_1 \text{ determines an edge of } \text{NE}(\bar{X}) \text{ [Kol96, Lemma II.4.12]}. \end{aligned} \quad (20)$$

On the other hand, $\delta(g_{n+1}) < 0 \Rightarrow (C_1, C_1) > 0$ (assertion 1), which implies that there exists $\beta \in \mathbb{Q}$ such that with respect to the basis $(C_1, C_2 + \beta C_1)$ of $N_1(\bar{X})$, the intersection form is of the form $x_1^2 - x_2^2$. [Kol96, Lemma II.4.12] then implies that C_1 is in the interior of $\text{NE}(\bar{X})$. The preceding sentence together with (20) implies assertions 2b and 2c. The first statement of assertion 2 follows from assertions 2a, 2b and 2c. \square

Proof of Theorem 1.4. W.l.o.g. we may (and will) assume that $\delta \neq \text{deg}$ and use the notations of Theorem 1.4'. Pick $f \in \mathbb{C}[x, y] \setminus \{0\}$ and let \bar{D}_f be the closure in \bar{X} of the curve D_f defined by f in \mathbb{C}^2 , so that $\bar{D}_f \sim \text{deg}(f)C_1 + \delta(f)C_2$. Consequently, $\delta(f) \geq 0$ for all $f \in \mathbb{C}[x, y] \setminus \{0\}$ iff $\text{NE}(\bar{X})$ is generated by C_1 and C_2 . Assertion 1 then follows from assertion 2 of Theorem 1.4'. \square

We now prove assertion 2. Proposition 2.11 implies that assertions 2b and 2b' are equivalent. Therefore, by assertion 1, it suffices to show that either 2a or 2b' implies that δ is positive on $\mathbb{C}[x, y] \setminus \mathbb{C}$. Now if 2a holds, then $(C_1, C_1) < 0$ (Theorem 1.4'). A criterion of Grauert (adapted to the case of normal surfaces in [Sak84, Theorem 1.2]) then implies that C_1 is *contractible*, i.e. there is a map $\pi : \bar{X} \rightarrow \bar{X}'$ of normal analytic surfaces such that $\pi(C_1)$ is a point and $\pi|_{\bar{X} \setminus C_1}$ is an isomorphism. In particular δ is the pole along the irreducible curve at infinity on the compactification \bar{X}' of $X := \mathbb{C}^2$, and consequently δ is positive on $\mathbb{C}[x, y] \setminus \mathbb{C}$, as required. Now assume 2b' holds. Then Theorem 1.4' implies that $(C_1, C_1) = 0$. Assume (to the contrary of our goal) that there exists $f \in \mathbb{C}[x, y] \setminus \mathbb{C}$ such that $\delta(f) = 0$. Then we have $(\bar{D}_f, C_1) = (\text{deg}(f)C_1, C_1) = 0$, so that $\bar{D}_f \cap C_1 = \emptyset$. Proposition 2.11 then implies that g_{n+1} is a polynomial, which contradicts 2b'. It follows that δ is positive on $\mathbb{C}[x, y] \setminus \mathbb{C}$, which completes the proof of assertion 2. \square

Proof of Theorem 1.8'. A straightforward computation using the entries of the intersection matrix \mathcal{I} from (19) shows that

$$(C_i^*, C_j) = \delta_{ij} \quad (21)$$

where δ_{ij} is the Kronecker delta. Since $(C_1^* + \epsilon C_2, C_2) < 0$ for all $\epsilon > 0$, and since l_2 is an edge of $\text{NE}(\bar{X})$, identity (21) immediately implies assertion 1. To complete the proof of Theorem 1.8', it suffices to prove the (\Leftarrow) direction of assertions 2 and 3. Now if $\delta(g_{n+1}) \geq 0$, then $\text{NE}(\bar{X})$ is generated by C_1 and C_2 (assertion 2 of Theorem 1.4'), so that (21) implies that l_2^* is also an edge of $\text{Nef}(\bar{X})$. This implies the (\Leftarrow) direction of assertion 2. On the other hand, if $\delta(g_{n+1}) < 0$, then C_1 is in the interior of $\text{NE}(\bar{X})$ (assertion 2c of Theorem 1.4'). Since $(C_2^*, C_1) = 0$, it follows that $C_2^* \notin \text{Nef}(\bar{X})$, which implies the (\Leftarrow) direction of assertion 3, as required to complete the proof of Theorem 1.8'. \square

⁵Even though [Kol96, Lemma II.4.12] is proved for only *non-singular* surfaces, its proof goes through for arbitrary normal surfaces using the intersection theory due to [Mum61].

Proof of Theorem 1.8. W.l.o.g. we may (and will) assume that $\delta \neq \deg$ and use the notations of Theorems 1.4' and 1.8'. Let ϕ_δ be as in Proposition 2.20. At first consider the case that $\phi_\delta = 0$. Then $n = 0$ and the key forms of δ are $g_0 = x$ and $g_1 = y$ (Example 2.7). On the other hand, (17) implies that (3) holds, so that Theorem 1.8 is true in this case. Therefore we may (and will) assume that $\phi_\delta \neq 0$, and divide the proof into separate cases depending on $\delta(g_{n+1})$.

Case 1: $\delta(g_{n+1}) \geq 0$. In this case C_2^* is on an edge of $\text{Nef}(\bar{X})$ (assertion 2 of Theorem 1.8'). Since any nef divisor is a limit of ample divisors and large multiples of ample divisors have global sections, it follows that there exists $f_1, f_2, \dots \in \mathbb{C}[x, y]$ such that $\bar{D}_{f_k} \sim r_k(C_1 + s_k C_2)$ for some $r_k, s_k \in \mathbb{Q}_{>0}$ such that $\lim_{k \rightarrow \infty} s_k = \frac{m_\delta \delta(g_{n+1})}{d_\delta}$ (where \bar{D}_{f_k} 's are defined as in the proof of Theorem 1.4). Identity (17) then implies that (3) holds with equality in this case.

Case 2: $\delta(g_{n+1}) < 0$. In this case C_1 is in the interior of $\text{NE}(\bar{X})$ (Theorem 1.4', assertion 2c). [Kol96, Lemma II.4.12] (adapted to the case of normal surfaces as in footnote 5) implies that $\text{NE}(\bar{X})$ has an edge of the form $\{r(C_1 - aC_2) : r \geq 0\}$ for some $a \in \mathbb{Q}_{>0}$, and moreover, there exists $r > 0$ such that $rC_1 - arC_2 \sim \bar{D}_g$ for some $g \in \mathbb{C}[x, y]$. Then $\deg(g) = r$ and $\delta(g) = -ar$. Pick $f \in \mathbb{C}[x, y] \setminus \mathbb{C}$. Since the 'other edge' of $\text{NE}(\bar{X})$ is spanned by C_2 (Theorem 1.4', assertion 2a), it follows that $\bar{D}_f \sim sC_2 + t(C_1 - aC_2)$ for some $s \in \mathbb{Q}_{\geq 0}$ and $t \in \mathbb{Q}_{>0}$, and therefore,

$$\frac{\delta(f)}{\deg(f)} = \frac{s - ta}{t} \geq -a = \frac{\delta(g)}{\deg(g)}.$$

It follows that

$$\inf \left\{ \frac{\delta(f)}{d_\delta \deg(f)} : f \in \mathbb{C}[x, y] \setminus \mathbb{C} \right\} = \frac{\delta(g)}{d_\delta \deg(g)}. \quad (22)$$

On the other hand, (17) implies that

$$\alpha(\delta) = \frac{\delta(g_{n+1})}{d_\delta \deg(g_{n+1})}.$$

Lemma 2.21 and Corollary 2.22 then imply that (3) holds with equality iff g_{n+1} is a polynomial.

The assertions of Theorem 1.8 now follow from the conclusions of the above 2 cases. \square

Proof of Corollary 1.10. We continue to assume that $\delta \neq \deg$ and use the notations of the proof of Theorem 1.8. Identify $\text{Nef}(\bar{X})$ with its image in \mathbb{R}^2 via the map $a_1 C_1 + a_2 C_2 \mapsto (a_1, a_2)$. Note that

- (A) The 'upper edge' of $\text{Nef}(\bar{X})$ is $l_1^* = \{r(1, d_\delta) : r \in \mathbb{R}_{\geq 0}\}$ (Theorem 1.8') and $l_1^* \subseteq C_\delta$ (since $(1, d_\delta) = (\deg(f), \delta(f))$, where f is a generic linear polynomial in (x, y)).
- (B) C_δ contains the 'lower edge' of $\text{Nef}(\bar{X})$ iff g_{n+1} is a polynomial iff g_k is a polynomial for all k , $0 \leq k \leq n + 1$ (follows from combining Theorem 1.8, Lemma 2.21 and Corollary 2.22).

Since $\text{Nef}(\bar{X})$ is a closed cone and since C_δ contains the *ample cone* of \bar{X} , the above observations imply assertion 1. For assertion 2 note that δ determines an analytic compactification of \mathbb{C}^2

$$\begin{aligned} &\text{iff } C_1 \text{ is contractible} \\ &\text{iff } (C_1, C_1) < 0 \text{ (by Grauert's criterion [Sak84, Theorem 1.2])} \\ &\text{iff } \delta(g_{n+1}) > 0 \text{ (Theorem 1.4', assertion 1).} \end{aligned}$$

Since the arguments of the proof of Theorem 1.8 show that $\delta(g_{n+1}) \leq 0$ iff the closure of C_δ contains the positive x -axis, this completes the proof of assertion 2. The equivalence of assertions 3a and 3b follows from assertion 1 and Theorem 2.12. Since 3c clearly implies 3b, it remains to show that 3b \Rightarrow S_δ is finitely generated. Since C_δ is a rational cone, 3b implies that $\bar{S}_\delta := C_\delta \cap \mathbb{Z}^2$ is finitely generated. Since \bar{S}_δ is *integral* over S_δ (i.e. for every $s \in \bar{S}_\delta$, there is a positive integer m such that $ms \in S_\delta$), it follows that S_δ is also finitely generated, as required to complete the proof of the corollary. \square

Example 3.1 (An example where (3) does not hold). Let δ be the semidegree on $\mathbb{C}(x, y)$ defined as follows:

$$\delta(f(x, y)) := \deg_x (f(x, y)|_{y=x^{-1}+\xi x^{-2}}) \quad \text{for all } f \in \mathbb{C}(x, y) \setminus \{0\},$$

where ξ is an indeterminate. Then the key forms of δ are $x, y, y - x^{-1}$. Moreover,

$$\begin{aligned} d_\delta &= \max\{\delta(x), \delta(y)\} = \max\{1, -1\} = 1, \\ m_\delta &= \gcd(\delta(x), \delta(y), \delta(y - x^{-1})) = \gcd(1, -1, -2) = 1, \end{aligned}$$

and therefore (17) implies that

$$\alpha(\delta) = \delta(y - x^{-1}) / \deg(y - x^{-1}) = -2. \quad (23)$$

Now consider the surface \bar{X} from Proposition 2.10. Then the matrix \mathcal{M} (from Proposition 2.10) and the intersection matrix \mathcal{I} of C_1 and C_2 are:

$$\mathcal{M} = \begin{pmatrix} 1 & 1 \\ 1 & -2 \end{pmatrix}, \quad \mathcal{I} = \mathcal{M}^{-1} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ 1 & -1 \end{pmatrix}. \quad (24)$$

In the notation of the proof of Theorem 1.4, $\bar{D}_y \sim \deg(y)C_1 + \delta(y)C_2 = C_1 - C_2$. It follows from (24) that $(C, C) = -1/3 < 0$, so that [Kol96, Lemma II.4.12] implies that C spans an edge of the cone of curves on \bar{X} , i.e. the polynomial g from Case 2 of the proof of Theorem 1.8 is y . It then follows from identities (22) and (23) that

$$\inf \left\{ \frac{\delta(f)}{d_\delta \deg(f)} : f \in \mathbb{C}[x, y] \setminus \mathbb{C} \right\} = \frac{\delta(y)}{d_\delta \deg(y)} = -1 > \alpha(\delta).$$

Example 3.2 (The *semigroup of values* does not distinguish semidegrees that determine algebraic compactifications of \mathbb{C}^2). Let δ be the semidegree on $\mathbb{C}(x, y)$ defined as follows:

$$\delta(f(x, y)) := 2 \deg_x \left(f(x, y)|_{y=x^{5/2}+x^{-1}+\xi x^{-3/2}} \right) \quad \text{for all } f \in \mathbb{C}(x, y) \setminus \{0\},$$

where ξ is an indeterminate. Then the key forms of δ are $x, y, y^2 - x^5, y^2 - x^5 - 2x^{-1}y$, with corresponding δ -values 2, 5, 3, 1 respectively. Since the δ -value of the last key polynomial is *positive*, it follows from the arguments of the proof of Corollary 1.10 that δ determines an analytic compactification of \mathbb{C}^2 . But the last key form of δ is *not* a polynomial, so that the compactification determined by δ is *non-algebraic* (Theorem 2.12). On the other hand, it follows from our computation of the values of δ and Corollary 2.22 that the semigroup of values of δ on polynomials is

$$N_\delta := \{\delta(f) : f \in \mathbb{C}[x, y]\} = \{2, 3, 4, \dots\}.$$

Now let δ' be the weighted degree on (x, y) -coordinates corresponding to weights 2 for x and 3 for y . Then δ' determines an *algebraic* compactification of \mathbb{C}^2 , namely the weighted projective surface $\mathbb{P}^2(1, 2, 3)$. But $N_\delta = N_{\delta'}$.

Example 3.3 ($\text{NE}(\tilde{X})$ or $\text{Nef}(\tilde{X})$ is not determined by purely numerical conditions if $\delta(g_{n+1}) < 0$). Let δ' be the semidegree on $\mathbb{C}(x, y)$ defined as follows:

$$\delta'(f(x, y)) := \deg_x(f(x, y)|_{y=\xi x^{-2}}) \quad \text{for all } f \in \mathbb{C}(x, y) \setminus \{0\},$$

where ξ is an indeterminate; in other words, δ' is the weighted degree on $\mathbb{C}[x, y]$ corresponding to weights 1 for x and -2 for y . Then the key forms of δ' are x, y . Moreover,

$$\begin{aligned} d_{\delta'} &= \max\{\delta'(x), \delta'(y)\} = \max\{1, -1\} = 1, \\ m_{\delta'} &= \gcd(\delta'(x), \delta'(y)) = \gcd(1, -2) = 1. \end{aligned}$$

Let \bar{X}' be the surface associated to δ' via the construction in Proposition 2.10. Then the matrix \mathcal{I} of curves C'_1 and C'_2 at infinity on \bar{X}' is identical to \mathcal{I} from (24), and it is straightforward to see that the weighted dual graphs of the curves at infinity (with respect to $\mathbb{C}^2 = \text{Spec}(\mathbb{C}[x, y])$) on the minimal resolutions \tilde{X} and \tilde{X}' of respectively \bar{X} and \bar{X}' are also identical - see Figure 1 (here E_0 (resp. E_3) corresponds to the strict transforms of C_1 (resp. C_2) in the case of \tilde{X} , and strict transforms of C'_1 (resp. C'_2) in the case of \tilde{X}').




Figure 1: Dual graph of curves at infinity on \tilde{X} and \tilde{X}'

On the other hand, if \bar{D}'_y is the closure of the x -axis in \bar{X}' , then $\bar{D}'_y \sim \deg(y)C'_1 + \delta'(y)C'_2 = C'_1 - 2C'_2$. Since $\bar{D}_y = C_1 - C_2$ determines an edge of $\text{NE}(\bar{X})$, it follows that $\text{NE}(\bar{X}) \not\cong \text{NE}(\bar{X}')$ (via the natural isomorphism $N_1(\bar{X}) \cong N_1(\bar{X}')$ given by the mapping $C_1 \mapsto C'_1, C_2 \mapsto C'_2$). Consequently, it follows that the cones of curves and nef cones of \tilde{X} and \tilde{X}' are also not isomorphic.

References

- [Abh78] Shreeram S. Abhyankar. On the semigroup of a meromorphic curve. I. In *Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977)*, pages 249–414, Tokyo, 1978. Kinokuniya Book Store. 4

- [Bäd01] Lucian Bădescu. *Algebraic surfaces*. Universitext. Springer-Verlag, New York, 2001. Translated from the 1981 Romanian original by Vladimir Maşek and revised by the author. 8
- [CPR05] Antonio Campillo, Olivier Piltant, and Ana J. Reguera. Cones of curves and of line bundles “at infinity”. *J. Algebra*, 293(2):513–542, 2005. 4
- [CPRL02] Antonio Campillo, Olivier Piltant, and Ana J. Reguera-López. Cones of curves and of line bundles on surfaces associated with curves having one place at infinity. *Proc. London Math. Soc. (3)*, 84(3):559–580, 2002. 5
- [FJ04] Charles Favre and Mattias Jonsson. *The valuative tree*, volume 1853 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2004. 7, 9
- [FJ07] Charles Favre and Mattias Jonsson. Eigenvaluations. *Ann. Sci. École Norm. Sup. (4)*, 40(2):309–349, 2007. 1, 3
- [Jon12] Mattias Jonsson. Dynamics on Berkovich spaces in low dimensions. <http://arxiv.org/abs/1201.1944>, 2012. 3
- [Kol96] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996. 12, 13, 14
- [Mac36] Saunders MacLane. A construction for absolute values in polynomial rings. *Trans. Amer. Math. Soc.*, 40(3):363–395, 1936. 1, 3
- [MN13] Pinaki Mondal and Tim Netzer. How fast do polynomials grow on semialgebraic sets? <http://arxiv.org/abs/1305.1215>, 2013. 2
- [Mon10a] Pinaki Mondal. Projective completions of affine varieties via degree-like functions. <http://arxiv.org/abs/1012.0835>, 2010. 2
- [Mon10b] Pinaki Mondal. Towards a Bezout-type theory of affine varieties. <http://hdl.handle.net/1807/24371>, March 2010. PhD Thesis. 2
- [Mon11] Pinaki Mondal. Analytic compactifications of \mathbb{C}^2 part I - curvettes at infinity. <http://arxiv.org/abs/1110.6905>, 2011. 8, 9, 10
- [Mon13a] Pinaki Mondal. An effective criterion for algebraic contractibility of rational curves. <http://arxiv.org/abs/1301.0126>, 2013. 3, 4, 7, 8, 9, 10
- [Mon13b] Pinaki Mondal. Mori dream surfaces associated with curves with one place at infinity. <http://arxiv.org/abs/1312.2168>, 2013. Preprint. 4
- [Mum61] David Mumford. The topology of normal singularities of an algebraic surface and a criterion for simplicity. *Inst. Hautes Études Sci. Publ. Math.*, (9):5–22, 1961. 12
- [Sak84] Fumio Sakai. Weil divisors on normal surfaces. *Duke Math. J.*, 51(4):877–887, 1984. 12, 14

- [SS94] Avinash Sathaye and Jon Stenerson. Plane polynomial curves. In *Algebraic geometry and its applications (West Lafayette, IN, 1990)*, pages 121–142. Springer, New York, 1994. [4](#)